



Phytobiotics in Poultry Industry as Growth Promoters, Antimicrobials and Immunomodulators – A Review

Wafaa A. Abd El-Ghany

¹*Poultry Diseases Department, Faculty of Veterinary Medicine, Cairo University, 1211, Giza, Egypt*

Corresponding author's E-mail: wafaa.ghany@yahoo.com; ORCID: 0000-0003-1686-3831

Received: 02 Nov. 2020

Accepted: 20 Dec. 2020

ABSTRACT

Due to the hazardous use of antimicrobials in poultry production sector, development of drug resistance have become a worldwide problem. Therefore, using biotic or natural products, such as phytobiotics (phytogenics or botanicals) have received a great attention as antibiotic substitutes. The use of phytobiotics or their constituents have been considered as a relatively new class of natural herbs that gained popularity and acceptability among poultry farmers. The incorporation of several types of phytobiotic additives in poultry feed have proved their ability to enhance the productive performance of broilers as well as layers. Moreover, phytobiotics presented great efficacy in counteracting intestinal pathogenic microorganism while maintaining the population of normal inhabitant beneficial microflora. Immunostimulatory effect on both humoral and cellular immunity as well as antioxidant properties were recorded as characters of phytobiotics. Therefore, this review article aimed to give a spotlight on the uses of different types of phytobiotics as poultry dietary additives to improve the productive parameters, reduce the pathogenic intestinal bacteria, and potentiate the immune response, especially after vaccination processes.

Keywords: Antimicrobial, Immunity, Performance, Plants, Poultry

INTRODUCTION

From several decades till now, poultry industry has been recognized as an important subsector of agricultural and veterinary fields due to increasing demand for meat and eggs as low cost protein sources. Continues feeding by sub-therapeutic levels of antimicrobials as growth promoter agents or antimicrobial compounds have had a negative impact on the balance of normal inhabitants of gut microflora, accumulation of antibiotic tissues residues as well as developing new strains of drug-resistant pathogenic bacteria (Castanon, 2007). Therefore, in 2006, the European Union Commission banned using of antibiotics in animal feeds as a growth promoter in different countries (European Union Commission, 2005).

Phytobiotics are also termed as phytogenics or botanicals. They are defined as natural, less toxic, and residue free plant-derived compounds that have been used as feed additives for livestock production (Wang et al., 2008). Phytobiotics are composed of natural bioactive components or substances of plant origin including terpenoids, alkaloids, glycosides and phenolics (Shad et

al., 2014). Phytobiotics could be classified as herbs from flowering, non-woody and non-persistent plants, botanicals or spices from non-leaf parts like seeds, fruits, bark or root, essential oils or extracts and oleoresins (Bote, 2004).

Numerous studies have been conducted to demonstrate the effect of phytobiotics as growth promoter feed additives like prebiotics and probiotics to enhance overall performance parameters as well as health conditions of poultry (Yasodha et al., 2019; Özbudak, 2019). Phytobiotics have been also used as antimicrobial, antiparasitic, anticoccidial as well as immunostimulant agents in poultry field (Manafi, 2015; Gilani et al., 2018; Hafeez et al., 2020). Phytogenic substances were extremely studied in different species of monogastrics (Gheisar and Kim, 2018), rabbits (Alagawany et al., 2018; Al-Sagheer et al., 2019) and fish (Naiel et al., 2019).

So, the purpose of current article review was to spotlight on phytogenic compounds that used in poultry field and their effects on the productive performance, antimicrobial activities and immuno-stimulatory properties.

Production parameters

Improved growth parameters were detected in birds fed on different kinds of herbs, polysaccharides or essential oils components (Yasodha *et al.*, 2019). The enhancement of the growth performance parameters after supplementation of phytobiotics may depend on the synergistic mechanism among their active molecular complex (Hussein *et al.*, 2020a). Phytobiotics could maintain or improve normal intestinal architecture, increase the villus length and consequently increase the surface of intestinal absorption (Tabatabaei, 2016). It has been demonstrated that phytobiotics are able to stimulate saliva production, secretion of digestible enzymes and bile production resulting in improving the performance and digestibility (Alloui *et al.*, 2014). Moreover, phytobiotics enhance the digestion and utilization of protein in the intestine (El-Gendi, 1996), decrease the gut pathogens (Kubkomawa *et al.*, 2013) and increase *Lactobacillus* spp. count (Windisch *et al.*, 2008). It has been observed that supplementation with herbal feed additives can alter the histological structure of the intestine and indicated elevation of the intestinal villi by deepening of its crypts (Murali *et al.*, 2012), increasing the dendritic cells absorption capacity in the intestinal lumen, stimulation of toll-like receptors and activation of epithelial to release the mucosal cytokines. Alcicek *et al.* (2004) assumed that feeding of broilers on phytobiotics stimulated the secretion of high amount of intestinal mucus and consequently reducing the pathogens adhesion and establishment of gut microbial eubiosis.

Supplementing broiler feed or water with essential oil mixtures of thymol and cinnamaldehyde (Tiihonen *et al.*, 2010), thymol and star anise (Kim *et al.*, 2016), clove and cinnamaldehyde (Chalghoumi *et al.*, 2013), coriander (Ghazanfari *et al.*, 2015; Hady *et al.*, 2016), oregano (Hashemipour *et al.*, 2014), a mixture of oregano, anise, and citrus peel (Abdelnaser *et al.*, 2019), carvacrol (Jamroz *et al.*, 2006), a blend of carvacrol, cinnamaldehyde and capsicum oleoresin (Bravo and Ionescu, 2008) and ginger extract (Olaiifa *et al.*, 2019) had been detected for improving all performance parameters including; feed intake, feed conversion rate and body weight. Moreover, supplementation of broilers diet with garlic (Elagib *et al.*, 2013), a mixture of garlic, mushroom and propolis (Daneshmand *et al.*, 2012), turmeric powder (Ahmadi, 2010), guggul resin (Iranparast *et al.*, 2014), dried ground leaves of stevia (Atteh *et al.*, 2008) and black cumin seeds (Khalaji *et al.*, 2011) presented the enhancement of broilers performance. Studies on the effects of herbal compounds on the production of broilers

Japanese quail's revealed improvement of all performances (Manafi *et al.*, 2016).

The laying hens represented improvement in egg production, eggshell strength and thickness as well as internal egg quality after treatment with mixture of plant extracts and essential oils (Bölükbaşı *et al.*, 2008; Radwan *et al.*, 2008; Kaya *et al.*, 2013).

Antimicrobial effect

It has been demonstrated that phytochemical compounds of phytobiotics have a strong antimicrobial activity against Gram-positive and Gram-negative bacteria either *in vivo* (Al-Kassie, 2010; Daka, 2013) or *in vitro* environment (Al-Mariri and Safi, 2014). Some phytochemical compounds as alkaloid inhibit DNA synthesis (Karou *et al.*, 2005) and form saponin complexes with the cell membrane (sterols) which leading to cells damage and collapse (Morrissey and Osbourn, 1999). The antimicrobial effect of essential oils could refer their ability to penetrate through the bacterial membrane (Helander *et al.*, 1998), their chemical structure (Frag *et al.*, 1989) and their aromaticity (Bowles and Miller, 1993).

Modulation of the gut microflora by phytochemical compounds plays an important role in maintaining host health (Tollba *et al.*, 2012). Several studies revealed that phytochemical compounds or their extracts reduce the population of intestinal pathogenic organisms and their metabolites, but increase the number of intestinal normal and helpful microflora which relieve the intestinal challenge and immune stress and consequently increase intestinal performance (Liu *et al.*, 2014). It has been speculated that organic acids of some phytobiotic feed additives may lower the intestinal pH that led to inhibiting the pathogenicity of local pathogens and lowering the level of their toxic products (Manafi *et al.*, 2016).

Antibacterial effect

Essential oils could reduce the growth of common intestinal poultry pathogens. Inoculation of thyme and cinnamon in the broiler ration reduces the total bacterial as well as coliform count in the intestinal tract (Karangiya *et al.*, 2016). An *in vitro* study revealed that flowers of *Calendula* presented growth inhibitory effects on *Escherichia coli* (*E. coli*) (Arora *et al.*, 2013). However, some reports indicated the inhibitory effects of dietary phytobiotics or plant extracts on *E. coli* activity *in vivo* (Abd El-Ghany and Ismail, 2014; Diaz-Sanchez *et al.*, 2015; Elmenawey *et al.*, 2019). Diet containing thyme essential oil helped for increasing in *Lactobacillus* and decreasing in *E. coli* counts in the intestine of Japanese

quail (Khaksar et al., 2012). Essential oils of oregano revealed antimicrobial properties on broiler carcasses through reduction of the total bacterial count especially *Salmonella* spp. (Aksit et al., 2006). Short chain fatty acids of phytobiotics revealed growth promoting effect on the intestinal beneficial microbiota as well as controlling the growth of *Salmonella enteritidis* (*S. enteritidis*) (Hansen et al., 1997). Quail's diet containing phytobiotic feed additives had significant ($P \leq 0.05$) increase in the number of *Lactobacilli*, decrease in *E. coli* population and inhibit the growth of *Salmonella* in the intestinal tract (Dorman and Deans, 2000).

The efficacy of a mixture of seven essential oils that inhibited the growth of *Clostridium perfringens* (*C. perfringens*) *in vitro* has been reported by Si et al. (2009). In the field studies of Mitsch et al. (2004), Siragusa et al. (2008), McReynolds et al. (2009), Abudabos et al. (2018), El-Sheikh et al. (2018) and Hussein et al. (2020b), different phytobiotic feed additives caused reduction of intestinal colonization and proliferation of *C. perfringens*, lesion score and mortalities as well as improvement in performance of broilers and their carcass quality.

The reduction in *C. perfringens* virulence after supplementation with essential oil compounds may be related to the stimulation of some digestible enzymes like trypsin which inactivates α toxin of type A and β toxin of type C strains of *C. perfringens*, stabilization of natural resident gut microflora like *Lactobacillus* spp. and consequently inhibition of the organism pathogenicity (Cho et al., 2014).

Moreover, essential oils of lemon, green tea and turmeric blend proved great efficacy in reducing the count of *S. enteritidis* and *Campylobacter jejuni* on the surface of chicken's carcass (Murali et al., 2012). The Eucalyptus volatile oils have been found to have the ability to relief broilers complicated respiratory distress caused by *Mycoplasma gallisepticum* (Abd El-Ghany, 2008).

Antifungal and detoxifying effect

Phytogetic compound have antifungal properties. Akgul and Kivanc (1988) reported about the inhibitory effects of some spices and oregano components on some foodborne fungi. Essential oil of marjoram reduced the *in vitro* growth of *Aspergillus flavus*, *A. niger*, *A. ochraceus* and *A. parasiticus* up to 89% (Deans and Waterman, 1993). The oil of lemon and orange can reduce the formation of *A. flavus* (Hasan et al., 2005). It has been demonstrated that *Allium sativum* has a great (60-80%) *in vitro* antifungal activity against *Aspergillus* and *Penicillium* spp. (Afzal et al., 2010). Onion, garlic and

ginger revealed antifungal activity against *A. flavus*, *A. niger* and *Cladosporium herbaru*, *Cinnamomum verum* and *Piper nigrum* which was studied with successful results by Tago et al. (2011). Phytobiotics like *Ocimum gratissimum*, *Cymbopogon citratus*, *Xylopi aethiopica*, *Monodora myristica*, *Syzygium aromaticum* proved their ability to inhibit the formation of non sorbic acid which is a precursor in the pathway of aflatoxin synthesis (Auwah, 1996). Neem extract represented inhibitory effects on biosynthesis of aflatoxins (B and G) (Bhatnagar et al., 1990). Also, neem leaf extract and its oil have been found to inhibit the growth and morphology of *Penicillium* spp. and consequently prevent the production of ochratoxin A (Bhatnagar et al., 1990).

Anticoccidial effect

The anticoccidial activity of some herbal plants have been documented (Willis et al., 2013). Phytogetic compounds have been found to reduce the severity of *Eimeria* spp. infection in broilers by alleviation of droppings score, intestinal lesions score and also reducing oocyst shedding (Zyan et al., 2017). Numerous phytobiotics as *Atemesia annua* (Allen et al., 1997), *Astragalus membranaceus* and *Sophora flavescens* (Youn and Noh, 2001), green tea (Jang et al., 2007), *Ageratum conyzoid* (Nweze and Obiwulu, 2009), *Musa paradisiaca* (Anosa and Okoro, 2010), olive tree (De Pablos et al., 2010), oregano essential oil (Tsinas et al., 2011), *Carica papaya* leaf extract (Nghonjuyi et al., 2015) and coconut oil (Hafeez et al., 2020) have indicated an excellent anticoccidial activity against different types of *Eimeria* spp. in birds.

Some essential oils presented similar drug efficacy in prevention and control of coccidiosis in broilers. The oregano oil and other mixtures of oils were similar to ionophores lasalocid (Giannenas et al., 2003) and monensin (Oviedo-Rondón et al., 2006), while a mixture of carvacrol, camphor, cineole and thymol was similar to salinomycin (Bozkurt et al., 2014) in terms of reduction of shedding and lesions of different *Eimeria* spp. in broilers.

Combined experimental infection of *C. perfringens* and *E. maxima* has been ameliorated after dietary treatment of three breeds of broilers with Capsicum, *Curcuma longa* oleoresins (Kim et al., 2015) and *Allium hookeri* root (Lee et al., 2018).

Immunomodulatory effect

In poultry production, reduction of infection as well as improvement of production by stimulation of the immune system after using phytogetic substances were

investigated previously (Zaki *et al.*, 2016). The immunomodulatory mechanism of phytoactive substances in poultry have been studied (Hashemi and Davoodi, 2012). Polysaccharide are very important immunoactive components of phytobiotics (Xue and Meng, 1996). Phytoactive compounds also induce their immunomodulatory effects through increasing immune cells proliferation, arising cytokines expression and elevation of antibody titers (Lee *et al.*, 2010; Park *et al.*, 2011; Pourhossein *et al.*, 2015). The immunogenicity of phytobiotics could be manifested as increasing macrophages, lymphocytes and natural killer cells activities as well as stimulation of interferon production (Hashemi and Davoodi, 2010; Kumar *et al.*, 2014). Plants contain flavonoids, vitamin C and carotenoids are able to enhance the immune system (Craig, 1999). Some herbs and species like garlic, echinacea and liquoric have immunostimulatory properties due to their composition of vitamin C, carotenoids and flavonoids as well as their abilities to stimulate macrophages, lymphocytes and natural killer cells activities and interferon production (Frankic *et al.*, 2009). Detecting *in vitro* immunostimulatory effect of dandelion, mustard and safflower either on lymphocytes and macrophages of chickens was performed by Lee *et al.* 2007. The results indicated inhibition of tumor cell growth, antioxidant effects, stimulation of lymphocyte proliferation and nitric oxide production by macrophages. The immunostimulant effect of some essential oils extracts of phytobiotics may be due to the presence of certain compounds that may bind to Immunoglobulin G (Ig G) receptors which led to stimulation of immune response (Ahmed *et al.*, 2013). In the study of Placha *et al.* (2014), inoculation of thymol in the broiler diet increased the trans-epithelial electrical resistance of duodenal mucosa.

It has been found that mushroom and plant polysaccharides have immunomodulatory effects in chickens infested with *Eimeria tenella* (Guo *et al.*, 2004).

Potential antiviral activity of plant seeds was recorded (Yaseen, 2003). Different types of herbs mix, spices, plant extracts and essential oils presented enhancement of immune response of birds (Huang *et al.*, 2007; Pourali *et al.*, 2010; Kavyani *et al.*, 2012; Abou-Elkhair *et al.*, 2014; Awaad *et al.*, 2016).

A significant ($P < 0.05$) elevation of antibody Enzyme-linked immunosorbent assay (ELISA) titer after vaccination with Newcastle Disease Virus (NDV) vaccine (Chowdhury *et al.*, 2018) was recorded after feeding on clove bud and cinnamon plant, while peppermint essential oil helped in significant ($P < 0.05$) rising in

Haemagglutination Inhibition (HI) antibody titer against Avian Influenza (AI) virus vaccines (H9N2) in broiler chicks (Sultan *et al.*, 2017). In addition, significant ($P < 0.05$) increase in HI titers against NDV vaccine and sheep red blood cells were observed in broiler chicks treated with Aloe vera gel (1%) in the drinking water for six weeks (Darabighane *et al.*, 2017)

Laying hens presented significant ($P > 0.05$) increase in ELISA titer after vaccination with ND, Infectious Bronchitis (IB) and Infectious Bursal Disease (IBD) vaccines (Özek *et al.*, 2011). Landy *et al.* (2011) observed that inoculation of broiler ration with neem powder for six weeks resulting in an elevation of HI antibody titer against AI vaccine, but not against ND vaccine. Barbour *et al.* (2008) evaluated the effect of using Eucalyptus and peppermint essential oils during vaccination of ND and IBD. The results indicated improved health conditions of the birds associated with elevation of ELISA titers against the used virus vaccines. Moreover, an increase in IgG and IgM ELISA titers has been observed in chickens fed on oregano essential oils for three weeks period (Malayoglu *et al.*, 2010). Recently, Abdelnaser *et al.* (2019) indicated that treatment with essential oils of oregano, anise, and citrus peel at level of 125 gm/ton induced positive effects on the immune response of *C. perfringens* challenged broilers after vaccination against ND, IB, AI and IBD viruses as well as increasing in relative spleen weight.

CONCLUSION

Using of phytobiotics in poultry diet as a feed additive and considering them as an antimicrobial substitute has become a very essential and critical issue currently. Phytobiotic compounds could replace antibiotic growth promoters to improve the productive performance of chickens, act as antibacterial, antifungal and antiparasitic agents as well as it has been considered as potential immunostimulants especially after routine vaccination programs of chickens. This review spotlight on the significant using of phytobiotics in poultry field and industry as natural antibiotic alternatives to avoid the emerged problem of antibiotic resistance,

Competing interests

The author have not declared any conflict of interest.

REFERENCES

- Abd El-Ghany WA (2008). Assessment of the efficacy of certain antibiotics and volatile oils for the treatment of induced yersinia infection in broiler chickens. Journal of the Egyptian

- Veterinary Medical Association, 69: 283-307. Available at: <https://scholar.cu.edu.eg/?q=wafaaabdelghany/publications/assessment-efficacy-certain-antibiotics-and-volatile-oils-treatment-ind>
- Abd El-Ghany WA and Ismail M (2014). Tackling experimental colisepticaemia in broiler chickens using phytobiotic essential oils and antibiotic alone or in combination. A mixture of essential oils of *Oreganum aetheroleum* is more effective compared with ciprofloxacin in the treatment of *E. coli* in broiler chickens. Iranian Journal of Veterinary Research, 15: 110-115. DOI: <https://www.dx.doi.org/10.22099/ijvr.2014.2341>
- Abdelnaser AHT, Moustafa MMA and Abd El-Haleem MI (2019). Effect of using synbiotics and essential oils on performance parameters and immune response of necrotic enteritis challenged broiler chicks. Benha Veterinary Medical Journal, 36: 373-380. DOI: <https://www.dx.doi.org/10.21608/BVMJ.2019.15094.1051>
- Abudabos AM, Alyemni AH, Dafalla YM and Khan RU (2018). The effect of phytochemicals on growth traits, blood biochemical and intestinal histology in broiler chickens exposed to *Clostridium perfringens* challenge. Journal of Applied Animal Research, 46: 691-695. DOI: <https://www.doi.org/10.1080/09712119.2017.1383258>
- Abou-Elkhair R, Ahmed HA and Selim S (2014). Effects of black pepper (*Piper nigrum*), turmeric powder (*Curcuma longa*) and coriander seeds (*Coriandrum sativum*) and their combinations as feed additives on growth performance, carcass traits, some blood parameters and humoral immune response of broiler chickens. Asian Australasian Journal of Animal Sciences, 27: 847-854. DOI: <https://www.dx.doi.org/10.5713/2Fajas.2013.13644>
- Afzal R, Mughal S M, Munir M, Sultana K, Qureshi R, Arshad M and Laghari MK (2010). Mycoflora associated with seeds of different sunflower cultivars and its management. Pakistan Journal of Botany, 42: 435-445. Available at: [http://www.pakbs.org/pjbot/PDFs/42\(1\)/PJB42\(1\)435.pdf](http://www.pakbs.org/pjbot/PDFs/42(1)/PJB42(1)435.pdf)
- Ahmed ST, Hossain ME, Kim GM, Hwang JA, Ji H and Yang CJ (2013). Effects of resveratrol and essential oils on growth performance immunity, digestibility and fecal microbial shedding in challenged piglets. Asian Australasian Journal of Animal Sciences, 26: 683-690. DOI: <https://www.dx.doi.org/10.5713/2Fajas.2012.12683>
- Ahmadi F (2010). Effect of turmeric (*Curcumin longa*) powder on performance, oxidative stress state and some of blood parameters in broilers fed on diets containing aflatoxin. Global Veterinaria, 5: 312-317. Available at: [http://www.idosi.org/gv/gv5\(6\)10/4.pdf](http://www.idosi.org/gv/gv5(6)10/4.pdf)
- Akgul A, and Kivanc M (1988). Inhibitory effects of selected Turkish spices and oregano components on some foodborne fungi. International Journal of Food Microbiology, 6: 263-268. DOI: [https://www.doi.org/10.1016/0168-1605\(88\)90019-0](https://www.doi.org/10.1016/0168-1605(88)90019-0)
- Aksit M, Goksoy E, Kok F, Ozdemir D and Ozdogan M (2006). The impacts of organic acid and essential oil supplementations to diets on the microbiological quality of chicken carcasses. Archiv fur Geflugelkunde, 70: 168-173. Available at: <https://www.european-poultry-science.com/The-impacts-of-organic-acid-and-essential-oil-supplementations-to-133diets-on-the-microbiological-quality-of-chickencarcasses,QUIEPTQyMTcxOTQmTUIEPT2MTAxNA.html>
- Alagawany M, Abdel-Hack ME, Al-Sagheer AA, Naiel MA, Saadeldin IM and Swelum AA (2018). Dietary cold pressed water cress and coconut oil mixture enhances growth performance, intestinal microbiota, antioxidant status, and immunity of growing rabbits. Animals, 8: 212. DOI: <https://www.dx.doi.org/10.3390/2Fani8110212>
- Alcicek A, Bozkurt M and Çabuk M (2004). The effect of a mixture of herbal essential oils, an organic acid or a probiotic on broiler performance. South African Journal of Animal Science 34: 217-222. Available at: <https://hdl.handle.net/10520/EJC94382>
- Al-Kassie GAM (2010). The effect of thyme and cinnamon on the microbial balance in gastro intestinal tract on broiler chicks. International Journal of Poultry Science, 9: 495-498. DOI: <http://www.dx.doi.org/10.3923/ijps.2010.495.498>
- Allen PC, Lydon J and Danforth H (1997). Effects of components of *Artemisia annua* on coccidia infections in chickens. Poultry Science, 76: 1156-1163. DOI: <https://www.doi.org/10.1093/ps/76.8.1156>
- Alloui N, Alloui MN, Agabou A (2014). Application of herbs and phytochemical feed additives in poultry production - a review. Global Journal Animal Science Research, 2: 234-243. Available at: <http://archives.gjasr.com/index.php/GJASR/article/view/57>
- Al-Mariri A and Safi M (2014). *In vitro* antibacterial activity of several plant extracts and oils against some gram-negative bacteria. Iranian Journal of Medical Sciences, 39: 36. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24453392>
- Al-Sagheer AA, Abdel-Hack ME, Alagawany M, Naiel MA, Mahgoub SA, Badr MM, Hussein EO, ALOWAIMER AN and Swelum AA (2019). Paulownia leaves as a new feed resource: Chemical composition and effects on growth, carcasses, digestibility, blood biochemistry, and intestinal bacterial populations of growing rabbits. Animals, 9: 95. DOI: <https://www.dx.doi.org/10.3390/2Fani9030095>
- Anosa GN and Okoro OJ (2010). Anticoccidial activity of the methanolic extract of *Musa Paradisiaca* root in chickens. Tropical Animal Health and Production, 43: 245-248. DOI: <http://www.doi.org/10.1007/s11250-010-9684-1>
- Arora D, Rani A and Sharma A (2013). A review on phytochemistry and ethnopharmacological aspects of genus *Calendula*. Pharmacognosy Reviews, 7: 179-187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24347926>
- Atteh JO, Onagbesan OM, Tona K, Decuypere E, Geuns JM and Buyse J (2008). Evaluation of supplementary stevia (*Stevia rebaudiana, bertonii*) leaves and stevioside in broiler diets: effects on feed intake, nutrient metabolism, blood parameters and growth performance. Journal of Animal Physiology and Animal Nutrition, 92: 640-649. DOI: <https://www.doi.org/10.1111/j.1439-0396.2007.00760.x>
- Awaad MHH, Afify MAA, Zoulfekar SA, Mohammed FF, Elmenawy MA and Hafez HM (2016). Modulating effect of peppermint and eucalyptus essential oils on vVND infected chickens. Pakistan Veterinary Journal, 36: 350-355. Available at: http://www.pvj.com.pk/pdf-files/36_3/350-355.pdf
- Awuah RT (1996). Possible utilization of plant products in grain storage. Proceedings of the Workshop on Mycotoxins in Food in Africa, Nov. 6-10, International Institute of Tropical Agriculture, Benin, p. 32. Available at: <http://www.academicjournals.org/AJB>
- Barbour EK, Yaghi RH, Shaib HA, Tayeb IT and Sleiman FT (2008). Evaluation of an essential oil in treatment of immunosuppressed coinfecting broilers. Eurasian Journal of Sustainable Agriculture, 2: 212-218. Available at: <http://www.aensi.org/aejsa/2008/212-218.pdf>
- Bhatnagar D, Zeringue HJ and McCormick SP (1990). Neem leaf extracts inhibit aflatoxin biosynthesis in *Aspergillus flavus* and *Aspergillus parasiticus*. Proceedings of the USDA Neem Workshop, April 16-17, Beltsville, Maryland, pp. 118-127. Available at: <https://agris.fao.org/agris-search/search.do?recordID=US9164967>
- Bölükbaşı ŞC, Erhan MK and Kaynar Ö (2008). The effect of feeding thyme, sage and rosemary oil on laying hen performance, cholesterol and some proteins ratio of egg yolk and *Escherichia coli* count in feces. Archiv fur Geflugelkunde, 72: 231-237. Available at: <https://www.european-poultry-science.com/The-effect-of-feeding-thyme-sage-and-rosemary-oil-on-laying-hen-performance-cholesterol-and-some-proteins-ratio-of-egg-yolk-and-span-classws-name-Escherichia-Colispan-cou,QUIE>
- Bote CJL (2004). Bioflavonoid's effects reach beyond productivity. Feed Mix, 12: 12-15.

- Bowles BL and Miller AJ (1993). Antibotulinal properties of selected aromatic and aliphatic aldehydes. *Journal of Food production*, 56: 788-794. DOI: <https://www.doi.org/10.4315/0362-028x-56.9.788>
- Bozkurt M, Aysul N, Küçükylmaz K, Aypak S, Ege G, Çatli A U, Aküit H, Çöven F, Seyrek K and Çınar M (2014). Efficacy of in-feed preparations of an anticoccidial, multienzyme, prebiotic, probiotic, and herbal essential oil mixture in healthy and *Eimeria* spp.-infected broilers. *Poultry Science*, 93: 389-399. DOI: <https://www.doi.org/10.3382/ps.2013-03368>
- Bravo D, and Ionescu C (2008). Meta-analysis of the effect of a mixture of carvacrol, cinnamaldehyde and capsicum oleoresin in broilers. *Poultry Science*, 87: 75. Available at: https://www.researchgate.net/publication/298327084_Metaanalysis_of_the_effect_of_a_mixture_of_carvacrol_cinnamaldehyde_and_capsicum_oleoresin_in_broilers
- Castanon JIR (2007). History of the use of antibiotic as growth promoters. *Journal of Poultry Science*, 86: 2466-2471. DOI: <https://www.doi.org/10.3382/ps.2007-00249>
- Chalghoumi R, Belgacem A, Trabelsi I, Bouatour Y and Bergaoui R (2013). Effect of dietary supplementation with probiotic or essential oils on growth performance of broiler chickens. *International Journal of Poultry Science*, 12: 538-344. DOI: <http://www.dx.doi.org/10.3923/ijps.2013.538.544>
- Cho JH, Kim HJ and Kim IH (2014). Effects of phytogetic feed additive on growth performance, digestibility, blood metabolites, intestinal microbiota, meat color and relative organ weight after oral challenge with *Clostridium perfringens* in broilers. *Livestock Science*, 160: 82-88. DOI: <https://www.doi.org/10.1016/j.livsci.2013.11.006>
- Chowdhury S, Mandala G, Patraa A, Kumara P, Samantab I, Pradhanc S, Chowdhurya A, Mandala G, Patraa A, Kumara P et al. (2018). Different essential oils in diets of broiler chickens: 2. Gut microbes and morphology, immune response, and some blood profile and antioxidant enzymes. *Animal Feed Science and Technology*, 236: 39-47. DOI: <https://www.doi.org/10.1016/j.anifeedsci.2017.12.003>
- Craig WJ (1999). Health-promoting properties of common herbs. *American Journal of Clinical Nutrition*, 70: 491-499. DOI: <https://www.doi.org/10.1093/ajcn/70.3.491s>
- Daka D (2013). Antibacterial effect of garlic (*Allium sativum*) on *Staphylococcus aureus*: An *in vitro* study. *African Journal of Biotechnology*, 10: 666-669. Available at: <http://www.academicjournals.org/AJB>
- Daneshmand A, Sadeghi GH and Karimi A (2012). The effects of a combination of garlic, oyster mushroom and propolis extract in comparison to antibiotic on growth performance, some blood parameters and nutrients digestibility of male broilers. *Brazilian Journal of Poultry Science*, 14: 141-147. DOI: <https://www.doi.org/10.1590/S1516-635X2012000200009>
- Darabighane B, Mahdavi A, Aghjehgheshlagh F, Zarei A, Kasapidou El and Nahashon S (2017). Effect of Aloe vera and vitamin E supplementation on the immune response of broilers. *Revista Colombiana de Ciencias Pecuarias*, 30: 159-164. DOI: <http://www.dx.doi.org/10.17533/udea.rccp.v30n2a07>
- Deans SG and Waterman PG (1993). Biological Activities of Volatile Oils. In: *Volatile Oil Crops: Their Biology, Biochemistry and Production*, Hay, R.K.M. and P.G. Waterman (Eds.). Longman, London, UK, pp. 97-111. Available at: <https://agris.fao.org/agris-search/search.do?recordID=GB9405744>
- De Pablos LM, dos Santos MF, Montero E, Garcia-Granados A, Parra A and Osuna A (2010). Anticoccidial activity of maslinic acid against infection with *Eimeria tenella* in chickens. *Parasitology Research*, 107: 601-604. DOI: <https://www.doi.org/10.1007/s00436-010-1901-3>
- Diaz-Sanchez S, D'Souza D, Biswas D and Hanning I (2015). Botanical alternatives to antibiotics for use in organic poultry production. *Poultry Science*, 94: 1419-1430. DOI: <https://www.doi.org/10.3382/ps/pev014>
- Dorman HJ and Deans SG (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, 88: 308-316. DOI: <https://www.doi.org/10.1046/j.1365-2672.2000.00969.x>
- Elagib HAA, El-Amin WIA, Elamin KM and Malik HEE (2013). Effect of dietary garlic (*Allium sativum*) supplementation as feed additive on broiler performance and blood profile. *Journal of Animal Science Advances*, 3: 58-64. Available at: [Link](http://www.ijvets.com/pdf-files/Volume-8-no-4-2019/294-299.pdf)
- El-Gendi GM (1996). Effect of feeding dietary herbal feed additives on productive and metabolic responses of broiler chicks. *Egyptian Poultry Science*, 16: 395-412. Available at: [https://www.scirp.org/\(S\(czeh2tfqyw2orz553k1w0r45\)\)/reference/ReferencesPapers.aspx?ReferenceID=1656208](https://www.scirp.org/(S(czeh2tfqyw2orz553k1w0r45))/reference/ReferencesPapers.aspx?ReferenceID=1656208)
- Elmenaway MA, Mohammed FA, Abdel-Alim GA and Abdel-Monef MHH (2019). The impact of essential oils blend on experimental colisepticemia in broiler chickens. *International Journal of Veterinary Science*, 8: 294-299. Available at: <http://www.ijvets.com/pdf-files/Volume-8-no-4-2019/294-299.pdf>
- El-Sheikh SM, Khairy MH, Eleiwa NZ, Abdalla OE and El-Monsef AGA (2018). Effect of sanguinarine phytoibiotic, sodium butyrate compared to ampicillin on controlling necrotic enteritis in broiler chickens. In *Proceedings of the Veterinary Medicine In-between Health & Economy (VMHE)*, Cairo, Egypt, 16-19 October 2018, Volume 55. DOI: <https://doi.org/10.26873/SVR-668-2018>
- Europe Union Commission (2005). Ban on antibiotics as growth promoters in animal feed enters into effect. Regulation 1831/2003/EC on additives for use in animal nutrition, replacing Directive 70/524/EEC on additives in feed- stuffs, Brussels, 22 December. Available at: https://ec.europa.eu/commission/presscorner/detail/en/IP_05_1687
- Farag RS, Badei AZMA, Hewedi FM and El-Baroty GSA (1989). Antioxidant activity of some spice essential oils on linoleic acid oxidation in aqueous media. *Journal of the American Oil Chemists' Society*, 66: 792-799. DOI: <https://www.doi.org/10.1007/BF02653670>
- Frankic T, Voljg M, Salobir J and Rezar V (2009). Use of herbs and spices and their extracts in animal nutrition. *Actaagricultuae Slovenica*, 92: 95-102. Available at: <http://aas.bf.uni-lj.si/zootehnika/94-2009/PDF/94-2009-2-95-102.pdf>
- Ghazanfari S, Mohammadi Z and Moradi AM (2015). Effects of coriander essential oil on the performance, blood characteristics, intestinal microbiota and histology of broilers. *Brazilian Journal of Poultry Science*, 17: 419-426. DOI: <https://www.doi.org/10.1590/1516-635X1704419-426>
- Gheisar MM and Kim IH (2018). Phytoibiotics in poultry and swine nutrition – a review. *Italian Journal of Animal Science*, 17: 92-99. DOI: <https://www.doi.org/10.1080/1828051X.2017.1350120>
- Giannenas I, Florou-Paneri P, Papazahariadou M, Christaki E, Botsoglou NA and Spais AB (2003). Effect of dietary supplementation with oregano essential oil on performance of broilers after experimental infection with *Eimeria tenella*. *Archives of Animal Nutrition*, 57: 99-106. DOI: <https://www.doi.org/10.1080/0003942031000107299>
- Gilani SMH, Zehra S, Hassan F, Galani S and Ashraf A (2018). Effect of natural growth promoters on immunity, and biochemical and haematological parameters of broiler chickens. *Tropical Journal of Pharmaceutical Research*, 17: 627-633. DOI: <https://www.doi.org/10.4314/tjpr.v17i4.9>
- Guo FC, Kwakkel RP, Williams BA, Parmentier HK, Li WK, Yang ZQ and Verstegen MW (2004). Effects of mushroom and herb polysaccharides on cellular and humoral immune responses of *Eimeria tenella*-infected chickens. *Poultry Science*, 83: 1124-1132. DOI: <https://www.doi.org/10.1093/ps/83.7.1124>
- Hady MM, M Zaki MM, Abd El-Ghany WA and Korany RMS (2016). Assessment of the broilers performance, gut healthiness and carcass characteristics in response to dietary inclusion of dried coriander, turmeric and thyme. *International Journal of Environmental and*

- Agriculture Research, 2: 153-159. Available at: <https://issuu.com/ijoear-environmentalandagriculture>
- Hafeez A, Ullah Z, Khan RU, Qudrat U and Naz S (2020). Effect of diet supplemented with coconut essential oil on performance and villus histomorphology in broiler exposed to avian coccidiosis. *Tropical Animal Health and Production*, 52: 2499-2504. DOI: <https://www.doi.org/10.1007/s11250-020-02279-6>
- Hansen LL, Larsen AE, Jensen BB and Hansen-Moller J (1997). Short time effect of zinc bacitracin and heavy fouling with faeces plus urine on boar taint. *Journal of Animal Science*, 64: 351-563. DOI: <https://www.doi.org/10.1017/S1357729800015927>
- Hasan MM, Chowdhury SP, Alam S, Hossain B and Alam MS (2005). Antifungal effects of plant extracts on seed-borne fungi of wheat seed regarding seed germination, Seedling health and vigour index. *Pakistan Journal of Biological Sciences*, 8: 1284-1289. DOI: <http://www.dx.doi.org/10.3923/pjbs.2005.1284.1289>
- Hashemi SR and Davoodi H (2010). Phytochemicals as new class of feed additive in poultry industry. *Journal of Animal and Veterinary Advances*, 9: 2295-2304. DOI: <http://www.dx.doi.org/10.3923/javaa.2010.2295.2304>
- Hashemi SR and Davoodi H (2012). Herbal plants as new immunostimulator in poultry industry: A review. *Asian Journal of Animal and Veterinary Advances*, 7: 105-116. DOI: <http://www.dx.doi.org/10.3923/ajava.2012.105.116>
- Hashemipour H, Kermanshahi H, Golian A and Khaksar V (2014). Effects of carboxy methyl cellulose and thymol + carvacrol on performance, digesta viscosity and some blood metabolites of broilers. *Journal of Animal Physiology and Animal Nutrition*, 98: 672-679. DOI: <https://www.doi.org/10.1111/jpn.12121>
- Helander IM, Alakomi HL, Latva-Kala K, Mattila-Sandholm T, Pol I, Smid EJ, Gorris LGM and von Wright A (1998). Characterization of the action of selected essential oil components on gram-negative bacteria. *Journal of Agricultural and Food Chemistry*, 46: 3590-3595. DOI: <https://www.doi.org/10.1021/jf980154m>
- Huang RL, Deng ZYC, Yang YL, Yin Xie MY, Wu GY, Li TJ, Li LL, Tang ZR, Kang P, Hou ZHP et al. (2007). Dietary oligochitosan supplementation enhances immune status of broilers. *Journal of Science of Food and Agriculture*, 87: 153-159. DOI: <https://www.doi.org/10.1002/jsfa.2694>
- Hussein EOS, Ahmed SH, Abudabos AM, Suliman GM, Abd El-Hack ME, Swelum AA and Alowaimer AN (2020a). Ameliorative effects of antibiotic-, probiotic- and phytochemical-supplemented diets on the performance, intestinal health, carcass traits, and meat quality of *Clostridium perfringens*-infected broilers. *Animals* 10: 669. DOI: <https://www.doi.org/10.3390/ani10040669>
- Hussein EOS, Ahmed SH, Abudabos AM, Aljumaah MR, Alkhlulaifi MM, Nassan MA, Suliman GM, Naiel MAE and Swelum AA (2020b). Effect of antibiotic, phytochemical and probiotic supplementation on growth, blood indices and intestine health in broiler chicks challenged with *Clostridium perfringens*. *Animals*, 10: 507. DOI: <https://www.doi.org/10.3390/ani10030507>
- Iranparast F, Parsaei S, Houshmand M and Naghiha A (2014). The effect of oral consumption of guggul (*Commiphora mukul*) resin on performance and humoral immunity response of broilers. *International Journal of Advanced Biological and Biomedical Research*, 2: 802-810. Available at: http://www.ijabbr.com/article_7237_c88f2b6f14b9d63e28ab48d317528e06.pdf
- Jamroz D, Wiertelcki T, Houszka M and Kamel C (2006). Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *Journal of Animal Physiology and Animal Nutrition*, 90: 255-268. DOI: <https://www.doi.org/10.1111/j.1439-0396.2005.00603.x>
- Jang SI, Jun MH, Lillehoj HS, Dalloul RA, Kong II-K, Kim S and Min W (2007). Anticoccidial effect of green tea-based diets against *Eimeria maxima*. *Veterinary Parasitology*, 144: 172-175. DOI: [https://www.doi.org/10.1016/s0304-4017\(01\)00385-5](https://www.doi.org/10.1016/s0304-4017(01)00385-5)
- Karangiya VK, Savsani HH, Patil SS, Garg DD, Murthi KS, Ribadiya NK and Vekariya SJ (2016). Effect of dietary supplementation of garlic, ginger and their combination on feed intake, growth performance and economics in commercial broilers. *Veterinary World*, 9: 245-250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27057106>
- Karou D, Savadogo A, Canini A, Yameogo S, Montesano C, Simporé J, Colizzi V and Traore A S (2005). Antibacterial activity of alkaloids from *Sida acuta*. *African Journal of Biotechnology*, 4: 1452-1457. Available at: <http://www.academicjournals.org/AJB>
- Kavyani A, Zare Shahne A, PorReza J, Jalali Haji-abadi SMA and Landy N (2012). Evaluation of dried powder of mushroom (*Agaricus bisporus*) as an antibiotic growth promoter substitution on performance, carcass traits and humoral immune responses in broiler chickens. *Journal of Medicinal Plants Research*, 6: 94-100. DOI: <https://www.doi.org/10.5897/JMPR11.1168>
- Kaya A, Kaya H, Macit M, Celebi S, Esenbuga N, Yoruk MA and Karaoglu M (2013). Effects of dietary inclusion of plant extract mixture and copper into layer diets on egg yield and quality, yolk cholesterol and fatty acid composition. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 19: 673-679. Available at: http://vetdergi.kafkas.edu.tr/extdocs/2013_4/673-679.pdf
- Khaksar V, Van Krimpen MM, Hashemipour H and Pilevar M (2012). Effects of thyme essential oil on performance, some blood parameters and ileal microflora of Japanese quail. *Journal of Poultry Science*, 49: 106-110. DOI: <https://www.doi.org/10.2141/jpsa.011089>
- Khalaji S, Zaghari M, Hatami K, Hedari-Dastjerdi S, Lotfi L and Nazarian H (2011). Black cumin seeds, *Artemisia leaves (Artemisia sieberi)*, and *Camellia L.* plant extract as phytochemical products in broiler diets and their effects on performance, blood constituents, immunity, and cecal microbial population. *Poultry Science*, 90: 2500-2510. DOI: <https://www.doi.org/10.3382/ps.2011-01393>
- Kim SJ, Lee KW, Kang CW and An BK (2016). Growth performance, relative meat and organ weights, cecal microflora, and blood characteristics in broiler chickens fed diets containing different nutrient density with or without essential oils. *Asian Australasian Journal of Animal Science*, 29: 549-554. DOI: <https://www.dx.doi.org/10.5713/2Fajas.15.0426>
- Kim JE, Lillehoj HS, Hong YH, Kim GB, Lee SH, Lillehoj EP and Bravo DM (2015). Dietary Capsicum and *Curcuma longa* oleoresins increase intestinal microbiome and necrotic enteritis in three commercial broiler breeds. *Research in Veterinary Science*, 102: 150-158. DOI: <https://www.doi.org/10.1016/j.rvsc.2015.07.022>
- Kubkomawa HI, Nafarnda DW, Mukang SM, Tizhe MA, Tuakam DK, Shua NJ, Ugwu CC, Opara MN, Neils JS and Okoli IC (2013). Ethno-Veterinary health management practices amongst livestock producers in Africa – A review. *World Journal of Agricultural Sciences*, 1: 252-257. Available at: <http://wsrjournals.org/journal/wjas>
- Kumar M, Kumar V, Roy D, Kushwaha R and Vaiswani S (2014). Application of herbal feed additives in animal nutrition – a review. *International Journal of Livestock Research* 4: 1-8. DOI: <https://www.dx.doi.org/10.5455/ijlr.20141205105218>
- Landy N, Ghalamkari G, Toghiani M and Yazdi F (2011). Humoral immune responses of broiler chickens fed with antibiotic and neem fruit powder (*Azadirachta Indica*) as feed additive supplemented diet. *International Conference of Life Science and Technology (IPCBE)*, 3: 153-155. Available at: <http://www.ipcbee.com/vol3/39-L20019.pdf>
- Lee YS, Lee SH, Gadde UD, Oh ST, Lee SJ and Lillehoj HS (2018). *Allium hookeri* supplementation improves intestinal immune response against necrotic enteritis in young broiler chickens. *Poultry Science*, 97: 1899-1908. DOI: <https://www.doi.org/10.3382/ps/pey031>

- Lee SH, Lillehoj HS, Chun HK, Tuo W, Park HJ, Cho SM, Lee YM and Lillehoj EP (2007). *In vitro* treatment of chicken peripheral blood lymphocytes, macrophages and tumor cells with extracts of Korean medicinal plants. *Nutrition Research*, 27: 362-366. DOI: <https://www.doi.org/10.1016/j.nutres.2007.04.001>
- Lee SH, Lillehoj HS, Hong YH, Jang SI, Lillehoj EP, Ionescu C, Mazuranok L and Bravo D (2010). *In vitro* effects of plant and mushroom extracts on immunological function of chicken lymphocytes and macrophages. *British Poultry Science*, 51: 213-221. DOI: <https://www.doi.org/10.1080/00071661003745844>
- Liu HN, Liu Y, Hu LL, Suo YL, Zhang L, Jin F, Feng XA, Teng N and Li Y (2014). Effects of dietary supplementation of quercetin on performance, egg quality, cecal microflora populations, and antioxidant status in laying hens. *Poultry Science*, 93: 347-353. DOI: <https://www.doi.org/10.3382/ps.2013-03225>
- Malayoglu HB, Baysal Ş, Misirlioğlu Z, Polat M, Yilmaz H and Turan N (2010). Effects of oregano essential oil with or without feed enzymes on growth performance, digestive enzyme, nutrient digestibility, lipid metabolism and immune response of broilers fed on wheat-soybean meal diets. *British Poultry Science*, 51: 67-80. DOI: <https://www.doi.org/10.1080/00071660903573702>
- Manafı M (2015). Comparison study of a natural non-antibiotic growth promoter and a commercial probiotic on growth performance, immune response and biochemical parameters of broiler chicks. *Journal of Poultry Science*, 52: 274-281. DOI: <https://www.doi.org/10.2141/jpsa.0150027>
- Manafı M, Hedayati M and Khalaji S (2016). Effectiveness of phyto-genic feed additive as alternative to bacitracin methylene disalicylate on hematological parameters, intestinal histomorphology and microbial population and production performance of Japanese quails. *Asian Australasian Journal of Animal Sciences*, 29: 1300-1308. DOI: <https://www.doi.org/10.5713/ajas.16.0108>
- McReynolds J, Waneck C, Byrd J, Genovese K, Duke S and Nisbet D (2009). Efficacy of multistrain direct-fed microbial and phyto-genetic products in reducing necrotic enteritis in commercial broilers. *Poultry Science*, 88: 2075-2080. DOI: <https://www.doi.org/10.3382/ps.2009-00106>
- Mitsch P, Zitterl-Eglseer K, Köhler B, Gabler C, Losa R and Zimpernik I (2004). The effect of two different blends of essential oil components on the proliferation of *Clostridium perfringens* in the intestines of broiler chickens. *Poultry Science*, 83: 669-675. DOI: <https://www.doi.org/10.1093/ps/83.4.669>
- Morrissey JP and Osbourn AE (1999). Fungal resistance to plant antibiotics as a mechanism of pathogenesis. *Microbiology and Molecular Biology Reviews*, 63: 708-724. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10477313>
- Murali N, Kumar-Phillips GS, Rath NC, Marcy J and Slavik MF (2012). Effect of marinating chicken meat with lemon, green tea and turmeric against food borne bacterial pathogens. *International Journal of Poultry Science*, 11: 326-332. DOI: <http://www.doi.org/10.3923/ijps.2012.326.332>
- Naiel MA, Ismael NE and Shehata SA (2019). Ameliorative effect of diets supplemented with rosemary (*Rosmarinus officinalis*) on aflatoxin B1 toxicity in terms of the performance, liver histopathology, immunity and antioxidant activity of Nile Tilapia (*Oreochromis niloticus*). *Aquaculture*, 511: 734264. DOI: <https://www.doi.org/10.1016/j.aquaculture.2019.734264>
- Nghonjuyi NW, Tiambo CK, Kimbi HK, Manka'a CN, Juliano RS and Lisita F (2015). Efficacy of ethanolic extract of *Carica papaya* leaves as a substitute of sulphamamide for the control of coccidiosis in Kabir chickens in Cameroon. *Journal of Animal Health and Production*, 3: 21-27. DOI: <http://www.doi.org/10.14737/journal.jahp/2015/3.1.21.27>
- Nweze NE and Obiwulu IS (2009). Anticoccidial effects of *Ageratum Conyzoides*. *Journal of Ethnopharmacology*, 25: 6-9 DOI: <https://www.doi.org/10.1016/j.jep.2008.11.014>
- Olaifa RO, Sogunle OM, Obileye LI, Ibitoye SA, Ayodeji TM, Odutayo OJ and Ogundele MA (2019). Effect of oral administration of three different phytobiotics on growth performance of locally-adapted turkeys. *EC Veterinary Science*, 4: 656-662. Available at: <https://www.econicon.com/ecve/pdf/ECVE-04-00162.pdf>
- Oviedo-Rondón EO, Hume ME, Hernández C and Clemente-Hernández S (2006). Intestinal microbial ecology of broilers vaccinated and challenged with mixed *Eimeria* species, and supplemented with essential oil blends. *Poultry Science*, 85: 8540860. DOI: <https://www.doi.org/10.1093/ps/85.5.854>
- Özbudak S (2019). Phytobiotics and their roles in broiler nutrition. *Journal of Poultry Research*, 16: 23-29. DOI: <https://www.doi.org/10.34233/jpr.465575>
- Özek K, Wellmann KT, Ertekin B and Tarım B (2011). Effects of dietary herbal essential oil mixture and organic acid preparation on laying traits, gastrointestinal tract characteristics, blood parameters and immune response of laying hens in a hot summer season. *Journal of Animal and Feed Sciences*, 20: 575-586. DOI: <https://www.doi.org/10.22358/jafs/66216/2011>
- Park IJ, Cha SY, Kang M, So YS, Go HG, Mun SP, Ryu KS and Jang HK (2011). Effect of proanthocyanidin-rich extract from *Pinus radiata* bark on immune response of specific-pathogen-free White Leghorn chickens. *Poultry Science*, 90: 977-982. DOI: <https://www.doi.org/10.3382/ps.2010-01160>
- Placha I, Takacova J, Ryzner M, Cobanova K, Laukova A, Strompfova V, Venglovska K and Faix S (2014). Effect of thyme essential oil and selenium on intestine integrity and antioxidant status of broilers. *British Poultry Science*, 55: 105-114. DOI: <https://www.doi.org/10.1080/00071668.2013.873772>
- Pourali M, Mirghelenj SA and Kermanshahi H (2010). Effects of garlic powder on productive performance and immune response of broiler chickens challenged with Newcastle disease virus. *Global Veterinaria*, 4: 616-621. Available at: [http://www.idosi.org/gv/gv4\(6\)10/16.pdf](http://www.idosi.org/gv/gv4(6)10/16.pdf)
- Pourhossein Z, Qotbi AAA, Seidavi A, Laudadio V, Centoducati G and Tufarelli V (2015). Effect of different levels of dietary sweet orange (*Citrus sinensis*) peel extract on humoral immune system responses in broiler chickens. *Animal Science Journal*, 86: 105-110. DOI: <https://www.doi.org/10.1111/asj.12250>
- Radwan N, Hassan RA, Qota EM and Fayek HM (2008). Effect of natural antioxidant on oxidative stability of eggs and productive and reproductive performance of laying hens. *International Journal of Poultry Science*, 7: 134-150. DOI: <http://www.doi.org/10.3923/ijps.2008.134.150>
- Shad AA, Ahmad S, Ullah R, AbdEl-Salam NM, Fouad H, Ur Rehman N, Hussain H and Saeed W (2014). Phytochemical and biological activities of four wild medicinal plants. *Scientific World Journal*, 2014: 857363. DOI: www.doi.org/10.1155/2014/857363
- Si W, Ni X, Gong J, Yu H, Tsao R, Han Y and Chambers JR (2009). Antimicrobial activity of essential oils and structurally related synthetic food additives towards *Clostridium perfringens*. *Journal of Applied Microbiology*, 106: 213-220. DOI: <https://www.doi.org/10.1111/j.1365-2672.2008.03994.x>
- Siragusa GR, Haas GJ, Matthews PD, Smith RJ, Buhr RJ, Dale NM and Wise MG (2008). Antimicrobial activity of lupulone against *Clostridium perfringens* in the chicken intestinal tract jejunum and caecum. *Journal of Antimicrobial Chemotherapy*, 61: 853-858. DOI: <https://www.doi.org/10.1093/jac/dkn024>
- Sultan R, Aslam A, Saleem G, Anjum A, Krull W, Kumosani T and Barbour B (2017). Studies on performance, immunity, and safety of broilers vaccinated with killed H9N2 vaccine and supplemented with essential oils of Mentofin® in drinking water. *International Journal of Applied Research and Veterinary Medicine*, 15: 76-84. Available at: <https://www.jarvm.com/articles/Vol15Iss2/Vol15%20iss2Barbour.pdf>

- Tabatabaei SN (2016). Effect of olibanum (*Boswellia thurifera*) as a feed additive on performance, some blood biochemical and intestinal morphology in broiler chicks. Research Opinions in Animal and Veterinary Sciences, 6: 130-134. Available at: <http://www.roavs.com/pdf-files/Issue-4-2016/130-134.pdf>
- Tagoe DNA, Nyarko HD and Akpaka R (2011). A comparison of the antifungal properties of onion (*Allium cepa*), ginger (*Zingiber officinale*) and garlic (*Allium sativum*) against *Aspergillus flavus*, *Aspergillus niger* and *Cladosporium herbarum*. Research Journal of Medicinal Plant, 5: 281-287. DOI: <http://www.dx.doi.org/10.3923/rjmp.2011.281.287>
- Tiihonen K, Kettunen H, Bento M H L, Saarinen M, Lahtinen S, Ouwehand AC, Schulze H and Rautonen N (2010). The effect of feeding essential oils on broiler performance and gut microbiota. British Poultry Science, 51: 381-392. DOI: <https://www.doi.org/10.1080/00071668.2010.496446>
- Tollba AAH, Shabaan SAM and Abdel-Mageed MAA (2012). Effects of using aromatic herbal extract and blended with organic acids on productive and physiological performance of poultry. Egyptian Poultry Science Journal, 30: 229-248. Available at: http://www.epsaegypt.com/pdf/2010_march/12-%201179.pdf
- Tsinas A, Giannenas I, Voidarou C, Tzora A and Skoufos J (2011). Effects of an oregano based dietary supplement on performance of broiler chickens experimentally infected with *Eimeria Acervulina* and *Eimeria Maxima*. Journal of Poultry Science, 48: 194-200. DOI: <https://www.doi.org/10.2141/jpsa.010123>
- Wang L, Piao XL, Kim SW, Piao XS, Shen YB and Lee HS (2008). Effects of *Forsythia suspensa* extract on growth performance, nutrient digestibility, and antioxidant activities in broiler chickens under high ambient temperature. Poultry Science, 87: 1287-1294. DOI: <https://www.doi.org/10.3382/ps.2008-00023>
- Willis WL, Wall DC, Isikhuemhen OS, Jackson JN, Ibrahim SA, Hur SL and Anike F (2013). Effect of level and type of mushroom on performance, blood parameters and natural coccidiosis infection in floor-reared broilers. Open Mycology Journal, 7: 1-6. DOI: <http://www.dx.doi.org/10.2174/1874437001307010001>
- Windisch W, Schedle K, Plitzner C and Kroismayr A (2008). Use of phytogenic products as feed additives for swine and poultry. Journal of Animal Science, 86: 140-148. DOI: <https://www.doi.org/10.2527/jas.2007-0459>
- Xue M, and Meng XS (1996). Review on research progress and prosperous of immune activities of bioactive polysaccharides. Journal of Traditional Veterinary Medicine, 3: 15-18. Available at: <http://www.jstage.jst.go.jp/browse/jpsa>
- Yaseen SA (2003). Potential antiviral activity of *Nigella sativa* extracts. Iraqi Journal of Veterinary Science, 17: 37-40. Available at: <https://eurekamag.com/research/004/276/004276713.php>
- Yasodha T, Jeevitha S, Yogesh S and Ramanan M (2019). Growth performance parameters of broiler chickens as influenced by herbal poultry feed. International Research Journal of Animal and Veterinary Sciences, 1: 26-30. Available at: <https://scirange.com/abstract/irjavs.2019.26.30>
- Youn HJ and Noh JW (2001). Screening of the anticoccidial effects of herb extracts against *Eimeria tenella*. Veterinary Parasitology, 96: 257-263. DOI: [https://www.doi.org/10.1016/s0304-4017\(01\)00385-5](https://www.doi.org/10.1016/s0304-4017(01)00385-5)
- Zaki MM, Abd El-Ghany WA, Hady MM and Korany RMS (2016). Effect of certain phytobiotics on the immune response of Newcastle disease vaccinated broiler chickens Asian Journal of Poultry Science, 10: 134-140. DOI: <http://www.dx.doi.org/10.3923/ajpsaj.2016.134.140>
- Zyan KA, Elshourbagy MA, Aggour G and Abdelfattah MA (2017). Molecular identification of *E. tenella* in broiler chicks in Kalyoubia governorate and evaluation of different strategies for control ceal coccidiosis. Benha Veterinary Medical Journal, 33: 175-182. DOI: <https://www.doi.org/10.21608/BVMJ.2017.43882>
- Windisch W, Schedle K, Plitzner C and Kroismayr A (2008). Use of phytogenic products as feed additives for swine and poultry. Journal of Animal Science, 86: 140-148. DOI: <https://www.doi.org/10.2527/jas.2007-0459>
- Xue M, and Meng XS (1996). Review on research progress and prosperous of immune activities of bioactive polysaccharides. Journal of Traditional Veterinary Medicine, 3: 15-18.
- Yaseen SA (2003). Potential antiviral activity of *Nigella sativa* extracts. Iraqi Journal of Veterinary Science, 17: 37-40.
- Yasodha T, Jeevitha S, Yogesh S and Ramanan M (2019). Growth performance parameters of broiler chickens as influenced by herbal poultry feed. International Research Journal of Animal and Veterinary Sciences, 1: 26-30. Available at: <https://scirange.com/abstract/irjavs.2019.26.30>
- Youn HJ and Noh JW (2001). Screening of the anticoccidial effects of herb extracts against *Eimeria tenella*. Veterinary Parasitology, 96: 257-263. DOI: [https://www.doi.org/10.1016/s0304-4017\(01\)00385-5](https://www.doi.org/10.1016/s0304-4017(01)00385-5)
- Zaki MM, Abd El-Ghany WA, Hady MM and Korany RMS (2016). Effect of certain phytobiotics on the immune response of Newcastle disease vaccinated broiler chickens Asian Journal of Poultry Science, 10: 134-140. DOI: <http://www.dx.doi.org/10.3923/ajpsaj.2016.134.140>
- Zyan KA, Elshourbagy MA, Aggour G and Abdelfattah MA (2017). Molecular identification of *E. tenella* in broiler chicks in Kalyoubia governorate and evaluation of different strategies for control ceal coccidiosis. Benha Veterinary Medical Journal, 33: 175-182. DOI: <https://www.doi.org/10.21608/BVMJ.2017.43882>