



## Effects of Broiler Breeders' Age on Egg Quality Characteristics and Their Correlation Coefficients

Freddy Manyeula<sup>1</sup>, Boingotlo Sebolai<sup>2</sup>, Godiraone Sempule<sup>1</sup>, and John Cassius Moreki<sup>1\*</sup>

<sup>1</sup>Department of Animal Science, Botswana University of Agriculture and Natural Resources, Private Bag 0027, Sebele, Gaborone, Botswana

<sup>2</sup>Department of Biometry and Mathematics, Faculty of Sciences, Botswana University of Agriculture and Natural Resources, Private Bag 0027, Sebele, Gaborone, Botswana

\*Corresponding author's Email: [jmoreki@buan.ac.bw](mailto:jmoreki@buan.ac.bw); ORCID: 0000-0003-2932-3359

Received: 21 July 2021

Accepted: 08 September 2021

### ABSTRACT

The current study was designed to assess the effect of Ross breeder hens' age on the egg qualities and their correlations. The external and internal qualities of eggs were compared, and their correlation coefficients as influenced by the age of breeder hens were determined. A sample of 300 Ross breeder hen eggs was obtained from the Ross breeder farm with 100 eggs drawn from each laying period of ages, namely 30, 45, and 60 weeks. Measured parameters included egg weight, egg length, egg width, shell weight, and shell thickness. Data were evaluated for linear and quadratic effects using polynomial contrasts. Response surface regression analysis was applied to describe the responses of egg quality to the aging of breeder hens. The results showed that egg weight, egg length, egg width, shell weight, egg yolk, egg content, egg volume, shell percentage, albumen weight, egg shape index, and egg surface area increased over time. Haugh unit and thick albumen indicated that the eggs in all age groups were fresh and had high quality. Shell thickness was constant in all age groups. Egg weight was significantly correlated with egg length, width, yolk (length, width, weight, and height), and shell weight. In conclusion, the egg quality improved as the hens' age increased implying that age is an effective factor in improving the quality of eggs.

**Keywords:** Age, Broiler breeder, Egg quality, Shell quality

### INTRODUCTION

An egg is a control house of nutrition for the growing embryo, and it is the source of essential amino acids and fatty acids for humans (Alkan et al., 2015). Measuring external and internal qualities offer an assurance of egg safety. At the farm level, the quality of the broiler breeder eggs is determined by egg external and internal qualities which in turn determine broiler chick weight at slaughter. Egg quality characteristics include cleanliness, freshness, egg weight, shell quality, yolk index, albumen index, Haugh unit, and chemical composition (Song et al., 2000; Roberts, 2004).

Quality depends on factors, such as time of oviposition, genotype, age, ambient temperature, and nutrition (Tumova and Gous, 2012). There are other less well-understood factors, including the effect of the breeder's age on egg quality before incubation, which may affect the embryonic life of the chick and thereafter the quality of the broiler chicks and growth potential post-

hatch. A vast quantity of literature has determined the effects of breeders' age on egg quality. Yilmaz and Bozkurt (2009) and Crosara et al. (2019) reported a significant deterioration in the shell characteristics as breeder hens aged. Kontecka et al. (2012) reported that as the reproductive season of the hens progressed egg weight increased while the percentage of the white (index and Haugh unit) decreases. Understanding the relationship between egg quality at different ages of broiler breeder hens is critical in production management. It is known that poor eggshell quality influences embryo development leading to low hatchability results (Kontecka et al., 2012). Nasri et al. (2020) observed that a positive correlation exists between the number of hatchlings and shell thickness and strength.

The internal and external egg quality characteristics in broiler breeder hens are largely unknown in Botswana because most research focuses on commercial layer chickens (Duman et al., 2016), indigenous chicken eggs

(Kgwatalala et al., 2016; Manyeula et al. 2018), and guinea fowl (Manyeula et al., 2020). It has been reported that poor egg quality causes high losses in egg production and high chick mortality in a broiler breeder production (Tona et al., 2004). Due to the lack of knowledge, some farmers would spend money on chicken feeds to improve the shell quality of aged chickens with low reproductive rates. Information on egg quality would be used to educate farmers on good management practices of raising breeder chickens to produce high-quality eggs. Hence, the objectives of this study were to evaluate the effect of broiler breeder hen's age on egg quality and to determine correlations among the different egg quality parameters at different ages. It was hypothesized that the age of breeder hens would affect the external and internal quality of eggs.

## MATERIALS AND METHODS

A total of 300 eggs (i.e., 100 eggs age period) were obtained from three commercial Ross broiler breeder hens aged 30, 45, and 60 weeks, which were reared in open-sided houses at Thamaga in Botswana. Eggs were transported to the Meat Science Laboratory at Botswana University of Agriculture and Natural Resources (BUAN), Botswana where they were stored overnight at 20°C and at 70-80% relative humidity (Ross Breeders, 2018). Eggs were used to study the effects of breeder age on egg quality characteristics and their correlations. Egg quality measurements were carried out the following day in the Meat Science Laboratory.

### Ethical approval

The experimental procedure employed conformed to the guideline for care and use of research animals and was approved by the Animal Research Ethics Committee of BUAN (AEC 2021-03).

### Management and diets

The feed, water, light program, and other management conditions were administered to broiler breeder hens in accordance with Ross breeder guidelines and recommendations (Ross Breeders, 2018). Vaccinations and medication were carried out following the company's comprehensive health management plan. Breeders were vaccinated by spray route against Newcastle disease and infectious bronchitis at 3, 6, 13, 18, and 24 weeks of age and thereafter at intervals of 8 weeks until breeders reached the end of their production life. Vaccinations against Newcastle disease were carried out

using Nobilis® ND Clone 30 and Nobilis® IB MA5, respectively. The vaccine against swollen head syndrome (SHS) was administered intramuscularly at 4, 14, and 24 weeks of age, whereas Panacur (active ingredient: Fenbendazole) was administered at 21 weeks and was repeated whenever signs of parasites were observed.

### Measurement of egg quality characteristics

A total of 300 eggs corresponding to each laying period, including weeks 30, 45, and 60 (100 eggs per laying period), were randomly sampled for egg quality analysis. At each period, the eggs were individually weighed using the OXO electronic scale (Explorer EX 224, OHAUS Corp, China) sensitive to 0.001 g. Egg length (EL) and egg width (EWD) were measured using a digital Vernier caliper (NEIKO 01407A Electronic Digital Caliper, 0-6 Inches, China) sensitive to 0.01 mm. Eggs were individually placed on a flat tile and cracked, and then egg yolk and albumen spread on the tile. Thereafter, the height of thick and thin albumen was measured using a Vernier caliper (0.01 mm). The egg yolk length (YL) and yolk width (YWD) were also measured using a Vernier caliper (0.01 mm). The egg yolk was then gently and smoothly transferred to a petri dish and weighed on an electronic food kitchen scale (0.001 g). The shell was then wiped with a paper cloth to remove adhering albumen and thereafter weighed using OXO electronic scale. The eggshell thickness (ST) was measured in three locations (broad, sharp, and equator) using a Vernier caliper (0.01 mm) and the values averaged. Egg content (EC), egg surface area (ESA), egg shape index (SI), Haugh unit (HU), egg volume (EV), shell percentage, and yolk index (YI) were calculated using the formulae given in Table 1.

### Statistical analysis

Data were evaluated for linear and quadratic effects using polynomial contrasts. Response surface regression analysis (SAS, 2010) was applied to describe the responses of egg quality to the aging of breeder hens using the quadratic model of  $y = ax^2 + bx + c$ , where  $y$  is the response variable,  $a$  and  $b$  signify the coefficients of the quadratic equation,  $c$  refers to intercept, and  $x$  stands for different breeder hen age (weeks). Correlation coefficients for egg weight and other egg quality traits were determined at different breeder ages using the Proc Corr procedure of SAS (2010) and tested for significance at a level of 0.05 ( $p \leq 0.05$ ).

**Table 1.** Formulae used to calculate some egg characteristics in the present study

Traits	Formula	References
Egg content weight (g)	Egg weight – shell weight	Alkan et al. (2015)
Shape index (%)	Egg weight / Egg length ×100	Anderson et al. (2004)
Haugh unit	100*log (albumen height+7.57-1.7*egg weight <sup>0.37</sup> )	Haugh (1937)
Egg volume (g/mm <sup>3</sup> )	0.708 × Egg weight × 100	Carter (1975)
Egg surface area (mm <sup>2</sup> )	3.978 × Egg weight <sup>0.7056</sup>	Alkan et al. (2015)
Shell percentage (%)	Shell weight / Egg weight × 100	Roberts (2004)
Yolk index (%)	Yolk height / Yolk width ×100	Duman et al. (2016)

**RESULTS**

**External parameters**

There were significant linear ( $p < 0.05$ ) trends on egg weight [ $y = 33.6 (\pm 4.51) - 0.81 (\pm 0.21) x$ ;  $R^2 = 0.61$ ;  $p < 0.05$ ] and length [ $y = 47.2 (\pm 2.49) - 0.30 (\pm 0.11) x$ ;  $R^2 = 0.42$ ;  $p < 0.05$ ] in response to breeder hen age (Table 2). The egg width [ $y = 33.7 (\pm 1.25) - 0.36 (\pm 0.06) x + 0.003 (\pm 0.0007) x^2$ ;  $R^2 = 0.51$ ;  $p < 0.05$ ] and shell weight [ $y = 8.6 (\pm 1.00) - 0.08 (\pm 0.05) x + 0.001 (\pm 0.0005) x^2$ ;  $R^2 = 0.05$ ;  $p < 0.05$ ] increased quadratically while shell percentage quadratically decreased at  $y = 84.4 (\pm 1.75) + 0.35 (\pm 0.08) x - 0.002 (\pm 0.0009) x^2$ ;  $R^2 = 0.61$ ;  $p < 0.05$  with the aging of breeder hens. However, no linear or quadratic trends ( $p > 0.05$ ) was observed in shell thickness as the breeder hens aged.

**Internal egg parameters**

Albumen weight [ $y = 26.3 (\pm 3.81) + 0.12 (\pm 0.18) x$ ;  $R^2 = 0.43$ ;  $p < 0.05$ ], egg surface area [ $y = 41.2 (\pm 3.56) - 0.66 (\pm 0.17) x$ ;  $R^2 = 0.61$ ;  $p < 0.05$ ], and egg volume [ $y = 26.8 (\pm 3.60) - 0.64 (\pm 0.17) x$ ;  $R^2 = 0.61$ ;  $p < 0.05$ ] linearly increased ( $p < 0.05$ ) in response to breeder hen age while shape index decreased linearly at  $y = 73.0 (\pm 3.40) + 0.20 (\pm 0.16) x$ ;  $R^2 = 0.05$ ;  $p < 0.05$  (Table 3). The thin albumen [ $y = 5.2 (\pm 0.91) + 0.06 (\pm 0.04) x - 0.0009 (\pm 0.0005) x^2$ ;  $R^2 = 0.17$ ;  $p < 0.05$ ], egg contents [ $y = 25.5 (\pm 4.19) + 0.89 (\pm 0.20) x - 0.005 (\pm 0.002) x^2$ ;  $R^2 = 0.63$ ;  $p < 0.05$ ] and Haugh unit [ $y = 63.3 (\pm 1.24) + 0.24 (\pm$

$0.06) x - 0.002 (\pm 0.0006) x^2$ ;  $R^2 = 0.30$ ;  $p < 0.05$ ] increased quadratically with breeder hen age. However, quadratic trends were observed only on the thick albumen [ $y = 5.2 (\pm 0.91) + 0.06 (\pm 0.04) x - 0.001 (\pm 0.0004) x^2$ ;  $R^2 = 0.17$ ;  $p < 0.05$ ]. There was a significant ( $p < 0.05$ ) quadratic decrease in yolk index [ $y = 34.4 (\pm 3.30) + 0.45 (\pm 0.15) x - 0.005 (\pm 0.002) x^2$ ;  $R^2 = 0.04$ ;  $p < 0.05$ ], whereas, yolk width [ $y = 25.2 (\pm 2.00) + 0.61 (\pm 0.09) x - 0.005 (\pm 0.002) x^2$ ;  $R^2 = 0.43$ ;  $p < 0.05$ ], height [ $y = 7.1 (\pm 1.07) + 0.46 (\pm 0.05) x - 0.005 (\pm 0.0006) x^2$ ;  $R^2 = 0.33$ ;  $p < 0.05$ ] and weight [ $y = -0.72 (\pm 1.65) + 0.77 (\pm 0.07) x - 0.007 (\pm 0.0009) x^2$ ;  $R^2 = 0.64$ ;  $p < 0.05$ ] increased quadratically in response to breeder hens' age (Table 4).

**Correlation coefficients at different ages**

There were significant correlations between egg weight and other egg quality traits at different breeder ages except for shell thickness and thin albumen (Table 5). Egg weight was positively correlated with egg length, egg width, yolk (length, width, weight, and height), and shell weight at different breeder ages. Thick albumin and egg weight correlated negatively at all ages. However, at 30 weeks the correlation was not significant ( $p > 0.05$ ) but significant ( $p < 0.05$ ) at 45 and 60 weeks of age (Table 5). Shell thickness was negative but not significantly correlated to egg weight at weeks 30 and 45, however, at week 60 a non-significant positive correlation coefficient was observed between egg weight and breeder hen age.

**Table 2.** Effect of Ross breeder hen age on the external egg parameters

Parameters	Age (week)				p value	
	30	45	60	SE	Linear	Quadratic
Egg weight (g)	54.1	61.6	67.2	0.43	< 0001	0.08
Egg length (mm)	54.8	57.7	59.6	0.23	< 0001	0.24
Egg width (mm)	41.8	43.9	44.7	0.11	< 0001	< 0001
Shell weight (g)	6.5	6.4	6.9	0.09	0.0007	0.03
Shell thickness (mm)	0.74	0.73	0.74	0.01	0.83	0.16
Shell percentage (%)	9.8	9.6	9.3	0.40	< 0001	0.02

SE: Standard error of the mean

**Table 4.** Effects of breeder age on yolk parameters of eggs in broiler breeder hens

Parameters	Age in weeks				p value	
	30	45	60	SE	Linear	Quadratic
Yolk index	43.13	43.94	42.37	0.31	0.08	0.002
Yolk width	38.70	41.79	42.44	0.19	< 0001	< 0001
Yolk height	16.64	18.33	17.96	0.10	< 0001	< 0001
Yolk weight	16.25	20.16	21.02	0.16	< 0001	< 0001

SE: Standard error of the mean

**Table 3.** Effect of age on internal egg quality in Ross breeder hens

Parameters	Age (week)				p value	
	30	45	60	SE	Linear	Quadratic
Albumen weight (g)	31.3	34.9	39.2	0.36	< 0001	0.44
Thin albumen (mm)	2.6	2.8	2.7	0.04	0.006	0.01
Thick albumen (mm)	6.1	5.8	5.1	0.08	< 0001	0.04
Shape index (%)	76.4	76.2	74.8	0.32	0.0006	0.12
Egg surface area (mm <sup>2</sup> )	58.0	63.9	68.3	0.36	< 0001	0.06
Egg contents (g)	47.6	55.1	60.1	0.34	< 0001	0.02
Egg volume (g/mm <sup>3</sup> )	43.1	49.1	53.6	0.34	< 0001	0.08
Haugh unit	68.8	70.2	70.9	0.11	< 0001	0.0001

SE: Standard error of the mean

**Table 5.** Correlation coefficients between egg weight and other egg quality traits at different ages of broiler breeder hens

Traits	30 weeks	45 weeks	60 weeks
Egg length	0.54*	0.70*	0.60*
Egg width	0.70*	0.64*	0.69*
Yolk length	0.33*	0.18*	0.47*
Yolk width	0.27*	0.23*	0.41*
Yolk weight	0.37*	0.56*	0.40*
Yolk height	0.27*	0.39*	0.31*
Thick albumin	-0.01 <sup>ns</sup>	-0.24*	-0.31*
Thin albumin	-0.08 <sup>ns</sup>	-0.06 <sup>ns</sup>	-0.16 <sup>ns</sup>
Shell weight	0.45*	0.36*	0.43*
Shell thickness	-0.29 <sup>ns</sup>	-0.05 <sup>ns</sup>	0.11 <sup>ns</sup>

<sup>ns</sup>: Not statistically significant at p < 0.05 level of significance, \*: Statistically significant at p < 0.05 level of significance

## DISCUSSION

### External egg parameters

Egg weight is an important phenotypic trait that influences egg quality and fitness of broiler breeder hens. It is known that egg weight increases with the breeder hen age (Kontecka et al., 2012). This increase in egg weight is related to the increase in weight of the yolk and albumin with a higher proportion of yolk than the proportion of albumin in the egg. This explains the reason for which egg weight (EWT), egg length, and EWD increased linearly with breeder hen age in the current study. These results are in line with several investigators who reported that EWT, EL, and EWD increase as broiler breeder hens grow older

(Luquetti et al., 2004; Abudabos 2010; Traldi et al., 2011). Similarly, Alsobayel (2013) observed that EWT of Cobb breeder was the lowest at 30-35 weeks and the highest at 40-45 weeks of age.

Good shell quality must be maintained throughout all the reproductive life of breeder hens since it influences embryonic development (Crosara et al., 2019). However, in the current study, as the breeder hens aged, SWT increased quadratically due to an increase in the egg content resulting in weak shell strength. It is known that shell weight and shell strength are negatively correlated (Tumova et al., 2014). Strong shell strength protects the egg from bacteria ingress. Vast literature supports the current results (Abudabos, 2010; Tumova et al., 2014;

Abudabos *et al.*, 2017). Shell thickness (ST) is one of the most important external quality parameters that affect shell breaking strength (Tumova *et al.*, 2014). In the current study, it was observed that ST was not affected by age of breeder hens suggesting that mineralization was constant. However, shell strength decreased with age because hens at an advanced age have low calcium reserves. Lack of linear and quadratic trends on ST could be due to the supplementation of calcium in the diet hence better shell quality across the ages. Similarly, van den Brand *et al.* (2004) reported no significant difference of ST across the age. Contrary to the current results, Padhi *et al.* (2013) reported higher ST at 52 and 72 weeks of age, compared to lower age. The current results imply that breeder age did not affect ST, indicating that shell quality was not affected. This could be attributed to the breeder hens being offered calcium supplements at the farm.

#### Internal parameters

The increased linear trends observed in albumen weight, SI, ESA, and SP with breeder hens' age could be attributable to EWT and body weight. These results compare favorably with those of Rath *et al.* (2015) and Carter (1975). Also, Manyeula *et al.* (2018) reported an increase in ESA with an increased body weight of indigenous Tswana hens. Similarly, Moreki *et al.* (2011) observed that ESA increased over time from 36 to 60 weeks of age in Ross broiler breeder hens. In the current study, a linear decrease of SI in response to breeder hens' age indicated that the young hens were likely to produce round eggs and older hens (60 weeks) normal-shaped eggs that were good for the setting. Egg shape index is the ratio of EWT to EL and is an important criterion for determining egg quality. Altuntaş and Şekeroğlu (2008) found that sharp, normal (standard), and round eggs have SI values of <72, 72-76, and >76, respectively. Similar results were reported by Roberts (2004) and Kontecka *et al.* (2012). Additionally, Nikolova and Kocevski (2006) found SI values of 76.16% and 74.20% in younger (45 weeks) and older (> 45 weeks) hens, respectively. Egg weight is influenced by breeder hen's age and this explains why thin albumen weight and EC quadratically increased with breeder hen's age (Suk and Park, 2001). It has been reported that HU is used to measure the quality of albumen (Rizzi and Chiericato, 2005) and freshness of eggs, thus high-quality eggs have thicker whites (Charvatova and Tumova, 2010). The study by Carrazzoni de Menezes *et al.* (2012) reported that the HU of fresh eggs decreases with age. The quadratic increase of HU in response to breeder hen age in the present study could be

due to good storage time and indicated that the egg was fresh (USDA, 2000). Indeed, Barbosa (2003) confirmed that good storage time and temperature caused an increase in HU and EWT. The present results are consistent with Alkan *et al.* (2015) who reported increased HU with increased egg weight. A decreased quadratic trend observed in thick albumen in response to breeder hen ages indicates that young chickens (30 weeks) have better thick albumen than the older flock (60 weeks).

Kontecka *et al.* (2012) reported that increased egg weight with age is due to the increasing weight of the yolk rather than white. This explains why yolk parameters quadratically increased with breeder hen age. Similar results were reported by Vieira and Moran (1998) and Padhi *et al.* (2013) who found increased yolk parameters with hen age. The proportion of yolk is positively related to egg size which is influenced by hen age (Rizzi and Chiericato, 2005; Johnston and Gous, 2007) supporting the results of the current study. Yolk index determines fresh and good quality eggs. Eggs with YI value ranging between 0.42 and 0.40 reflects good quality egg (Sharp and Powell, 1930). The YI values in this study were within the range of 0.40-0.42 implying that as hens get older the quality of eggs also improves. Thus, age could be a factor that determines the quality of broiler breeder hen eggs.

#### Correlation coefficients at different ages

The EWT showed a strong, positive significant correlation with EL, EWD, and YL in all weeks signifying that selection for EWT will automatically lead to improvement in EL and EWD regardless of the hen's age. Previous studies by Apuno *et al.* (2011), Manyeula *et al.* (2020), Obike and Azu (2012), Tebesi *et al.* (2012), and Alkan *et al.* (2015) also revealed positive strong significant correlations between the EWT and EL. These results were expected because EL and EWD are important factors determining egg weight. The significant positive correlation between EWT and YL, YWD, Yolk weight (YWT), and YH could be attributable to the fact that egg yolk occupies egg width area, hence contributing to heavier egg (egg weight). The YWT constitutes the yolk portion which may have influenced YWD and YL. However, the strong positive and significant correlations between EWT and YWT in all investigated weeks in the current study are consistent with those reported by Padhi and Rajkumar (2013) and Alkan *et al.* (2015). This indicates that grading heavy eggs for hatching will lead to a great improvement in yolk parameters which is a source of energy and lipids for the development of the embryo. Speaker (1998) confirmed that energy and lipids for



embryo development are stored in the yolk sac. The thick albumen height was negatively correlated to EWT at weeks 45 and 60 implying that an increase in EWT corresponds to a decrease in albumen height. However, this result was not expected since albumen is a very important component contributing to EWT. The present result contradicts reports by Ukwu et al. (2017) and Abdalla (2018). The differences found between results may be partially attributable to differences in strains/lines, nutrition, and environment. The regression analysis between the age and shell weight found a negative quadratic trend suggesting that the younger hens had heavier shells, compared to older hens. Such results were expected since the younger breeder hens (30 weeks) have more calcium reserves compared to breeder hens aged 45-80 weeks in a study by Suk and Park (2001). Similarly, Abdalla (2018) reported a positive significant association between egg weight and shell weight. Roberts (2004) states that the eggshell thickness is affected by nutrition, stress, disease, and production system. However, the correlation between shell thickness and egg weight was not significant at all breeder hen ages (Table 5), suggesting that mineralization was not affected by the aging of the breeder hens. This could be due to the addition of supplementation of minerals in their diets. These findings are in consonance with Aryee et al. (2020) who found a non-significant correlation between egg weight and shell weight.

## CONCLUSION

The findings of the study period led to the conclusion that Ross breeder hens' age determines internal and external quality traits. Also, egg quality is affected by egg weight as indicated by positive correlation values. The present results suggest that the selection of breeder eggs according to breeder age will simultaneously lead to improvement in other egg quality parameters.

## DECLARATIONS

### Acknowledgments

The authors sincerely acknowledge Ross Breeders Botswana for donating eggs used in this study. Moreover, the authors express their gratitude to the Botswana University of Agriculture and Natural Resources for granting the use of laboratory facilities for analysis.

### Competing interests

The authors declare no conflict of interest.

## Authors' contributions

Dr. F. Manyeula conceptualized this study and together with Mr. G. Sempule carried out the investigation. Drs Manyeula and B. Sebolai and Mr. Sempule were responsible for data curation. Drs Manyeula, Sebolai, and JC Moreki wrote, edited, and reviewed the manuscript. Dr. Moreki also served as the corresponding author and together with Dr. Manyeula worked on the suggestions made by the reviewers.

## Ethical considerations

Prior to submission, the authors checked ethical issues including plagiarism, consent to publish, misconduct, data fabrication, and double publication.

## REFERENCES

- Abdalla AM (2018). Effect of laying hens age on some egg quality traits with emphasis on their correlation. *International Journal of Livestock Research*, 8: 76-82. DOI: <http://www.dx.doi.org/10.5455/ijlr.20180228055101>
- Abudabos A (2010). The effect of broiler breeder strain and parent flock age on hatchability and fertile hatchability. *International Journal of Poultry Science*, 9: 231-235. DOI: <http://www.dx.doi.org/10.3923/ijps.2010.231.235>
- Abudabos AM, Aljumaah RS, Algawaan AS, Al-Sornokh H, and Al-Atiyat RM (2017). Effects of hen age and egg weight class on the hatchability of free-range Indigenous chicken eggs. *Brazilian Journal of Poultry Science*, 19: 33-40. DOI: <https://www.doi.org/10.1590/1806-9061-2016-0264>
- Alkan S, Galic A, Karsli T, and Karabag K (2015). Effects of egg weight traits in partridge (*Alectoris chukar*). *Journal of Applied Animal Research*, 43(4): 450-456. DOI: <https://www.doi.org/10.1080/09712119.2014.980419>
- Alsobayel AA, Almarshade MA, and Albadry MA (2013). Effect of breed, age and storage period on egg weight, egg weight loss and chick weight of commercial broiler breeders raised in Saudi Arabia. *Journal of the Saudi Society of Agricultural Sciences*, 12: 53-57. DOI: <http://www.dx.doi.org/10.1016/j.jssas.2012.06.003A>
- Altuntaş E, and Şekeroğlu A (2008). Effect of egg shape index on mechanical properties of chicken eggs. *Journal of Food Engineering*, 85: 606-612. DOI: <http://www.dx.doi.org/10.1016/j.jfoodeng.2007.08.022>
- Anderson KE, Tharrington JB, Curtis PA, and Jones FT (2004). Shell characteristics of eggs from historic strains of single comb white leghorn chickens and relationship of egg shape to shell strength. *Introduction Journal of Poultry Science*, 3: 317-319. DOI: <http://www.dx.doi.org/10.3923/ijps.2004.17.19>
- Apuno AA, Mbap ST, and Ibrahim T (2011). Characterization of local chickens (*Gallus gallus domesticus*) in shelleng and song local government areas of Adamawa State, Nigeria. *Agriculture and Biology Journal of North America*, 2: 6-14. Available at: <https://scihub.org/ABJNA/PDF/2011/1/ABJNA-2-1-6-14.pdf>
- Aryee G, Adu-Aboagye G, Shiburah ME, Nkrumah T, and Amedorme D (2020). Correlation between egg weight and egg characteristics in Japanese Quail. *International Journal of Animal Science and Technology*, 8(3): 51-54. DOI: <http://www.dx.doi.org/10.11648/j.avs.20200803.11>
- Carrazzoni de Menezes P, Rodrigues de Lima E, Pinto de Medeiros J, Ketry de Oliveira WN, and Evêncio-Neto J (2012). Egg quality of

- laying hens in different conditions of storage, ages and housing densities. *Revista Brasileira de Zootecnia*, 41(9): 2064-2069. DOI: <https://www.doi.org/10.1590/S1516-35982012000900014>
- Carter TC (1975). The hen's egg. Estimation of shell superficial area and egg volume using fresh measurements of egg weight and shell length and breadth alone or in combination. *Brazilian Poultry Science*, 16: 541-543. DOI: <https://www.doi.org/10.1080/00071667508416224>
- Charvátová V, and Tůmová E (2010). Time of oviposition and egg composition: A review. *Scientia Agriculturae Bohemica*, 41: 190-195. Available at: <https://agris.fao.org/agris-search/search.do?recordID=CZ2011000104>
- Crosara FSG, Pereira VJ, Lellis CG, Barra KC, Santos SKA, Souza LCGM, Morais TA, Litz FH, Limão VA, Braga PFS *et al.* (2019). Is the eggshell quality influenced by the egg weight or the breeder age? *Brazilian Journal of Poultry Science*, 21: 1-8. DOI: <https://www.doi.org/10.1590/1806-9061-2018-0896>
- Duman M, Şekeroğlu A, Yildirim A, Elroglu H, and Camci O (2016). Relation between egg shape index and egg quality characteristics. *European Poultry Science*, 80: 1-9. DOI: <https://www.doi.org/10.1399/eps.2016.117>
- Gualhanone A, Furlan RL, Fernandez-Alarcon MF, and Macari M (2012). Effect of breeder age on eggshell thickness, surface temperature, hatchability and chick weight. *Brazilian Journal of Poultry Science*, 14: 9-14. DOI: <https://www.doi.org/10.1590/S1516-635X2012000100002>
- Haugh H (1937). The Haugh unit for measuring egg quality. *The U.S. Egg and Poultry Magazine*, 43: 552-555. DOI: <https://www.doi.org/10.3382/ps.0411461>
- Johnston SA, and Gous RM (2007). Modelling the changes in the proportion of the egg components during a laying cycle. *British Poultry Science*, 48: 347-353. DOI: <https://www.doi.org/10.1080/00071660701381134>
- Kgwatalala PM, Molapisi M, Thutwa K, Sekgopi B, Selemoge TP, and Nsoso SJ (2016). Egg quality characteristics and phenotypical correlations among egg quality traits in the naked neck, normal and dwarf strains of Tswana chickens raised under intensive management system. *International Journal of Environmental and Agriculture Research*, 2: 96-104. Available at: <https://ijear.com/Paper-August-2016/IJOEAR-APR-2016-39.pdf>
- Kontecka H, Nowczewski S, Sierszula MM, and Witkiewicz K (2012). Analysis of changes in egg quality of broiler breeders during the first reproduction period. *Anniversary Animal Science*, 12: 609-620. DOI: <http://www.dx.doi.org/10.2478/v10220-012-0051-1>
- Luquetti BC, Gonzales E, Bruno LDG, Furlann RL, and Macari M (2004). Egg traits and physiological neonatal chick parameters from broiler breeder at different ages. *Brazilian Journal of Poultry Science*, 6: 13-17. DOI: <https://www.doi.org/10.1590/S1516-635X2004000100002>
- Manyeula F, Tsopito CM, Kamau JM, Mogotsi M, Nsoso SJ, and Moreki JC (2018). Effect of *Imbrasia belina* (Westwood) or *Vigna subterranea* (L) Verde or *Tylosema esculentum* (Burchell) Schreiber as protein sources in diets fed to Tswana hens on egg quality. *Journal of Animal Science and Veterinary Medicine*, 3: 190-196. DOI: <http://www.dx.doi.org/10.31248/JASVM2018.118>
- Manyeula F, Tumagole O, and Kgwatalala PM (2020). Phenotypic correlations among various egg quality traits in pearl grey, Lavender, Royal purple, and white varieties of helmeted guinea fowl. *Journal of World's Poultry Research*, 10: 580-586. DOI: <http://www.dx.doi.org/10.36380/jwpr.2020.66>
- Moreki JC, van Der Merwe HJ, and Hayes JP (2011). Effect of dietary calcium level on egg production and eggshell quality in broiler breeder hens from 36 to 60 weeks of age. *Online Journal of Animal and Feed Research*, 1(1): 01-07. Available at: <http://www.ojafr.ir/main/attachments/article/54/OJAFR,%20A%2001.pdf>
- Nasri H, van den Brand H, Najjar T, and Brouzouaia M (2020). Egg storage and breeder age impact on egg quality and embryo development. *Journal of Animal Physiology and Animal Nutrition*, 104: 257-268. DOI: <http://www.dx.doi.org/10.1111/jpn.13240>
- Nikolova N, and Kocevski D (2006). Forming egg shape index as influenced by ambient temperatures and age of hens. *Biotechnology in Animal Husbandry*, 22: 119-125. DOI: <http://www.dx.doi.org/10.2298/BAH0602119N>
- Obike OM, and Azu KE (2012). Phenotypic correlations among body weight, external and internal egg quality traits of pearl and black strains of guinea fowl in a humid tropical environment. *Journal of Animal Science Advance*, 2: 857-864. Available at: <https://www.semanticscholar.org/paper/Phenotypic-Correlations-among-Body-Weight%2C-External-Obike-Azu/b918090339a6f0b9fdee120ee007c42b04c91fb0>
- Padhi MK, Chatterjee RN, Haunshi S and Rajkumar U (2013). Effect of age on egg quality in chicken. *Indian Journal of Poultry Science*, 4(48): 122-125. Available at: <https://krishi.icar.gov.in/jspui/bitstream/123456789/4159/1/MKP%20IJPS%20EQ.pdf>
- Rath PK, Mishra PK, Mallick BK, and Behura NC (2015). Evaluation of different egg quality traits and interpretation of their mode of inheritance in White Leghorns. *Veterinary World*, 8: 449-452. DOI: <http://www.dx.doi.org/10.14202/vetworld.2015.449-452>
- Rizzi C, and Chiericato GM (2005). Organic farming production. Effect of age on the productive yield and egg quality of hens of two commercial hybrid lines and two local breeds. *Italian Journal of Animal Science*, 4: 160-162. DOI: <https://www.doi.org/10.4081/ijas.2005.3s.160>
- Roberts J (2004). Factors affecting egg internal quality and eggshell quality in laying hens. *Journal of Poultry Science*, 41: 161-177. Available at: <https://agris.fao.org/agris-search/search.do?recordID=JP2005002175>
- Roberts JR, Chousalkar K, and Samiullah A (2013). Egg quality and age of laying hens: Implications for product safety. *Animal Production Science*, 53: 1291-1297. DOI: <https://www.doi.org/10.1071/AN12345>
- Ross Breeders (2018). Parent Stock Management Handbook. [https://en.aviagen.com/assets/Tech\\_Center/Ross\\_PS/RossPSHandBook2018.pdf](https://en.aviagen.com/assets/Tech_Center/Ross_PS/RossPSHandBook2018.pdf)
- Sharp PF, and Powel CK (1930). Decrease in the interior quality of hen's eggs during storage as indicated by the yolk. *Industrial and Engineering Chemistry*, 22: 908-910. DOI: <https://www.doi.org/10.1021/ie50248a031>
- Song KT, Choi SH, and Oh HR (2000). A comparison of egg quality of pheasant, chukar, quail and guinea fowl. *Asian-Australasian Journal of Animal Science*, 7: 986-990. DOI: <https://www.doi.org/10.5713/ajas.2000.986>
- Suk YO, and Park C (2001). Effect of breed and age of hens on the yolk to albumen ratio in two different genetic stocks. *Poultry Science*, 80: 855-858. DOI: <https://doi.org/10.1093/ps/80.7.855>
- Tebesi T, Madibela OR, and Moreki JC (2012). Effect of storage time on internal and external characteristics of Guinea fowl (*Numida meleagris*) eggs. *Journal of Animal Science in Advance*, 2: 534-542. Available at: <https://www.semanticscholar.org/paper/Effect-of-Storage-Time-on-Internal-and-External-of-Tebesi-Madibela/1b94f58dc2ab492b783ed3e84dd11170d932c07a>
- Tona K, Onagbesan O, De Ketelaere B, Decuyper E, and Bruggeman V (2014). Effects of age of broiler breeders and egg storage on egg quality, hatchability, chick quality, chick weight, and chick posthatch growth to forty-two days. *Journal of Applied Poultry Research*, 13: 10-18. DOI: <https://www.doi.org/10.1093/japr/13.1.10>
- Traldi AB, Menten JFM, Silva CS, Rizzo PV, Pereira PWZ, and Antaroza J (2011). What determines hatchling weight: breeder age

- or incubated egg weight? *Brazilian Journal of Poultry Science*, 3: 283-285. DOI: <https://www.doi.org/10.1590/S1516-635X2011000400011>
- Tumová E, and Gous RM (2012). Interaction between oviposition time, age, and environmental temperature and egg quality traits in laying hens and broiler breeders. *Journal of Animal Science*, 57: 451-459. DOI: <http://www.dx.doi.org/10.17221/6411-CJAS>
- Tumová E, Gous RM, and Tyler N (2014). Effect of hen age, environmental temperature, and oviposition time on eggshell quality and eggshell and serum mineral contents in laying and broiler breeder hens. *Czech Journal of Animal Science*, 59(9): 435-443. DOI: <http://www.dx.doi.org/10.17221/7655-CJAS>
- Ukwu HO, Abari PO, and Kuusu DJ (2017). Principal component analysis of egg quality characteristics of Isa brown layer chickens in Nigeria. *World Scientific News*, 70(2): 304-311. Available at: <http://www.worldscientificnews.com/wp-content/uploads/2017/01/WSN-702-2017-304-311.pdf>
- United States Department of Agriculture (USDA) (2000). Egg-grading manual. Washington Department of Agriculture 56p. Agricultural Marketing Service, p. 75. Available at: <https://naldc.nal.usda.gov/download/CAT11094176/PDF>
- Van den Brand H, Parmentier HK, and Kemp B (2005). Effect of housing system (outdoor vs. cages) and age of laying hens on egg characteristics. *British Poultry Science*, 45(6): 745-752. DOI: <http://www.dx.doi.org/10.1080/00071660400014283>
- Vieira SL, and Morann JRT (1998). Eggs and chicks from broiler breeders of extremely different age. *Journal of Applied Poultry Research*, 7: 372-376. DOI: <https://www.doi.org/10.1093/japr/7.4.372>
- Yilmaz AA, and Bozkurt Z (2009). Effects of hen age, storage period and stretch film packing on internal and external quality traits of table egg. *Lucrări științifice Zootehnieși Biotehnologii*, 42(2): 1-8. Available at: <https://www.semanticscholar.org/paper/EFFECTS-OF-HEN-AGE%2C-STORAGE-PERIOD-AND-STRETCH-FILM-Yilmaz-Bozkurt/7bc77baa4ff7c74496d3e3c1759d1f1de4feddb7>