



# Effects of Date Seed Flour on Broiler Chickens' Growth Performance, Apparent Digestibility of Protein, and Apparent Metabolizable Energy

Ilmiatus Sholichatunnisa<sup>1</sup> , Osfar Sjojfan<sup>2</sup> , Tri Eko Susilorini<sup>2</sup> , Danung Nur Adli<sup>2</sup> , and Muhammad Halim Natsir<sup>2\*</sup> 

<sup>1</sup>Postgraduate Student of Faculty of Animal Science, University of Brawijaya, Malang, East Java, Indonesia

<sup>2</sup>Faculty of Animal Science, University of Brawijaya, Malang, East Java, Indonesia

Corresponding author's E-mail: [emhanatsir@ub.ac.id](mailto:emhanatsir@ub.ac.id)

Received: 27 June 2022  
Accepted: 14 August 2022

## ABSTRACT

The use of agricultural by-products as alternative feed ingredients is recommended to reduce production costs and maximize income. This study aimed to determine the effects of added date seed flour on broiler chickens' growth performance, protein digestibility, and metabolic energy. A total of 200 Lohmann MB-202 day-old chicks were randomly allocated to 5 treatments and 4 replication with 10 broiler chickens per cage. The treatments used in the current research included control without the addition of date seed flour (T0), basal feed + 2.5% date seed flour (T1), basal feed + 5.0% date seed flour (T2), basal feed + 7.5% date seed flour (T3), basal feed + 10% date seed flour (T4). The investigated parameters were growth performance, apparent digestibility of protein, and apparent metabolizable energy-nitrogen (AMEn). The result showed that adding date seed flour significantly affected final body weight, apparent digestibility of protein, and AMEn. In contrast, the date seed flour was no significant effect on the feed intake feed conversion ratio and income over feed cost. In conclusion, the addition of 10% date seed flour successfully increases final body weight, apparent digestibility of protein, and AMEn without any adverse effect on the broiler chickens.

**Keywords:** Broiler chicken, Date seed flour, Metabolizable energy, Performance, Protein digestibility

## INTRODUCTION

Intensive poultry production systems are increasingly demanding a supply of protein. Generally, the success of poultry rearing depends on breed, feed, and management. Feed costs contributed to approximately 65-75% of poultry industry production (Sjojfan et al., 2021a). As farmers face the problem of the rising price of feed, it is crucial to use potentially available feed sources (Adli, 2021a). This case would be reducing the cost of poultry feeds as well as simultaneously contributing to reducing foot carbon print (Adli, 2021b). One potentially raw feed that can be used for broiler chickens is date palm (*Phoenix dactylifera* L.).

Dates are quite famous in Muslim countries, including Indonesia, as well as middle-east countries (Risa et al., 2018). In the Middle East, dates are the main staple food. In addition, the date palm is relatively adaptable to tropical areas, compared to subtropical areas. Thus, this

condition would be advantageous for tropical countries like Indonesia. Dates consisted of minerals, such as iron, potassium, selenium, calcium, and vitamins (C, B<sub>1</sub>, and B<sub>2</sub>) (Primurdia and Kusnadi, 2014). The glucose content, crude protein, and crude fiber of date palms are around 50-57%, 1.8-2%, and 2-4%, respectively (Kresnadipayana and Lestari, 2017). Tareen et al. (2017) reported 7.5% fat and 5.8% protein content for date seeds. The amino acids consisted of 0.17% methionine, 0.31% lysine, and 0.36% threonine, which is higher than maize (Tareen et al., 2017).

The nutritional content of dates depends on their strain and level of maturity (Giovanny et al., 2019). Although dates are rich in nutritional content, the seeds should be processed to increase shelf-life, palatability, and digestibility. Accordingly, this study aimed to determine the effects of added date seed flour on growth performance, protein digestibility, and metabolic energy of broiler chickens

## MATERIALS AND METHODS

### Ethical approval

Ethical approval for the study was given by the Animal Care and Use Committee, University of Brawijaya, Indonesia No. 44-KEP-UB-2022.

### Experimental design

A total of 200 MB-202 Lohmann chicks aged one day old (JAPFA Comfeed Ltd. Commercially) were randomly allocated to 5 treatments and 4 replicates of 10 broilers per cage. Each cage was adjusted with installed nipple drinks, free access water (*ad libitum*), rice hulls (130 × 100 cm, × 60 cm), and a controlled environment. The feed was given twice a day at the morning (07.00 a.m.) and afternoon (4 p.m.).

The treatments used in the research were the control without the addition of date seed flour (T0), basal feed + 2.5% date seed flour (T1), basal feed + 5.0% date seed flour (T2), basal feed + 7.5% date seed flour (T3), and basal feed + 10% date seed flour (T4). The investigated parameters included growth performance, apparent digestibility of protein, apparent metabolic energy (AME), and apparent metabolizable energy-nitrogen corrected (AMEn). The formulated feed (starter periods) consisted of maize, soybean meal, meat bone meal, corn gluten meal, crude palm oil, custom mineral mix, custom premix, and date seed (BSN, 2015a; Table 1). The finisher diet consisted of maize, soybean meal, meat bone meal, rice bran, crude palm oil, custom mineral mix, custom premix, and date seed (BSN, 2015b; Table 2).

### Preparation of the date seed flour

The current study was conducted on dates (*Phoenix dactylifera* L.). The preparation step of making date seed flour was the following (Bouaziz et al., 2021). The mean weight of the dates was 25 g. First, the dates were subjected to reverse osmosis (RO) water. Afterward, dates were dried using the oven at 60°C and ground into a powder, followed by 3-day storage at room temperatures. Before beginning the trial, dates were taken for analyses of proximate (Sjofjan et al., 2021b).

### Growth performance

The broiler chickens' weights were recorded at the beginning and the end of the weeks. In line, feed intake was measured as the differences between the amount of feed given and the remaining feed. Income over feed cost (IOFC) was calculated by considering final body weight

multiples with broiler price at the site and differences between feed intake and feed price (Setyawan et al., 2019). In the end, the feed conversion ratio (FCR) was expressed by dividing the amount of feed given by the total weight gained (Adli, 2021b).

### Apparent digestibility of protein and apparent metabolizable energy-nitrogen corrected

At the end of the experiments (35 days old), a total of 20 broiler chickens were removed into metabolic-artificial cages. The feed intake of the broiler chickens was recorded for apparent digestibility of protein on day 37 and AMEn on day 38.

Both AME and AMEn were measured according to the following formulae:

$$AME = AME - (ANR/FI)$$

$$AMEn = AME - (8.22 \times ANR/FI)$$

Where, ANR is apparent Nitrogen retention, FI denotes feed intake. The correction factor is 8.22 (Sjofjan et al., 2021a).

The apparent digestibility of protein is calculated using the following formulae.

$$\text{Protein digestibility} = (A \times B) - (C \times D)$$

$$(A \times B)$$

Where, A is feed consumption, B signifies protein in feed, C accounts for the amount of excreta, and D denotes protein (%) in excreta (Regar and Kowel, (2021).

### Statistical analysis

For the statistical analysis, analysis of variance (ANOVA) using a general linear model (GLM) was carried out using SAS OnDemand for Academics (ODA, Cary, NC, USA). The results were presented as standard error mean (SEM). Moreover, probability values were calculated using the least significant different testing. The following model was used:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where,  $Y_{ij}$  signifies the parameters observed,  $\mu$  is the overall mean,  $T_i$  denotes the effect level of date seed flour, and  $e_{ij}$  is the amount of error number. The treatments included control without the addition of date seed flour (T0), basal feed + 2.5% date seed flour (T1), basal feed + 5.0% date seed flour (T2), basal feed + 7.5% date seed flour (T3), basal feed + 10% date seed flour (T4). P value less than 0.05 was considered statistically significant. Moreover, probability values were calculated using the least significant different testing if there differ significantly ( $p < 0.05$ ).

**Table 1.** Composition of broiler chicken diet at starter period

Ingredients (% as is basis)	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
Yellow maize	43.55	41.50	38.55	36.05	33.55
Maize gluten meal	22.45	22.45	22.45	22.45	22.45
Soya bean meal	25.00	25.00	25.00	25.00	25.00
Meat and bone meal 50	2.50	2.50	2.50	2.50	2.50
Poultry meat meal	2.50	2.50	2.50	2.50	2.50
Palm olein	1.00	1.00	1.00	1.00	1.00
Salt	0.15	0.15	0.15	0.15	0.15
Custom Mineral mix*	1.00	1.00	1.00	1.00	1.00
Custom Vitamin mix**	1.00	1.00	1.00	1.00	1.00
Date seed flour	-	2.50	5.00	7.50	10.00
Total	100	100	100	100	100
<b>Calculated composition</b>					
ME (Kcal/kg)	2982.00	2969.88	2957.77	2945.65	2933.54
Crude Protein (CP)	21.56	21.56	21.56	21.56	21.56
Crude fibre (CF)	3.60	3.78	3.97	4.15	4.34
Calcium (Ca)	3.0	3.0	3.05	3.11	3.05
Phosphorus (P)	1.5	1.5	1.5	1.5	1.5
Lysine	1.21	1.21	1.21	1.21	1.21
Methionine	0.70	0.70	0.70	0.70	0.70
<b>Proximate composition</b> (Wet Chemical analyses)					
ME (Kcal/kg)	2833	2873	2832	2811	2833
Crude Protein (CP)	21.01	21.00	21.00	21.00	21.00
Crude fibre (CF)	4.00	4.00	4.00	4.00	4.00
Calcium (Ca)	1.2	1.2	1.2	1.2	1.2
Phosphorus (P)	1.75	1.75	1.75	1.75	1.75

\*: 12.5 mg Iron, 3 mg Copper, 37.5 mg Manganese, 31.32 mg Zinc, 5 mg Iodine, and 0.0625 mg Selenium; \*\*: 6000 IU Vitamin A, 1000 IU Vitamin D3, 10 mg Vitamin E, 1.5 mg Vitamin K3, 5 mg Vitamin B1, 2.5 mg Vitamin B2, 0.5 mg Vitamin B6, 2 mg Vitamin B12, 5.5 mg Niacin, 0.2 mg Pantothenic acid, 30 mg Betaine; \* T<sub>0</sub>: Control without the addition of date seed flour, T<sub>1</sub>: Basal feed + 2.5% date seed flour, T<sub>2</sub>: Basal feed + 5.0% date seed flour, T<sub>3</sub>: Basal feed + 7.5% date seed flour, T<sub>4</sub>: Basal feed + 10% date seed flour

**Table 2.** Composition of broiler chicken diet at finisher period

Ingredients (% as is basis)	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
Yellow maize	43.40	41.50	38.55	36.05	33.55
Maize gluten meal	19.45	19.45	19.45	19.45	19.45
Soya bean meal	25.00	25.00	25.00	25.00	25.00
Rice bran	9.00	9.09	9.09	9.09	9.09
Palm olein	1.00	1.00	1.00	1.00	1.00
Salt	0.15	0.15	0.15	0.15	0.15
Custom Mineral mix*	1.00	1.00	1.00	1.00	1.00
Custom Vitamin mix**	1.00	1.00	1.00	1.00	1.00
Date seed flour	-	2.50	5.00	7.50	10.00
Total	100	100	100	100	100
<b>Calculated composition</b>					
ME (Kcal/kg)	2983.70	2971.10	2971.10	2971.10	2971.10
Crude Protein (CP)	19.89	19.89	19.89	19.89	19.89
Crude fibre (CF)	4.30	4.49	4.49	4.49	4.49
Calcium (Ca)	2.0	2.0	2.0	2.0	2.0
Phosphorus (P)	0.5	0.5	0.5	0.5	0.5
Lysine	0.8	0.8	0.8	0.8	0.8
Methionine	0.3	0.3	0.3	0.3	0.3
<b>Proximate composition</b> (Wet Chemical analyses)					
ME (Kcal/kg)	2900	2900	2900	2900	2900
Crude Protein (CP)	19.00	19.00	19.00	19.00	19.00
Crude fibre (CF)	4.00	4.00	4.00	4.00	4.00
Calcium (Ca)	1.5	1.5	1.5	1.5	1.5
Phosphorus (P)	0.5	0.5	0.5	0.5	0.5

\*: 12.5 mg Iron, 3 mg Copper, 37.5 mg Manganese, 31.32 mg Zinc, 5 mg Iodine, and 0.0625 mg Selenium; \*\*: 6000 IU Vitamin A, 1000 IU Vitamin D3, 10 mg Vitamin E, 1.5 mg Vitamin K3, 5 mg Vitamin B1, 2.5 mg Vitamin B2, 0.5 mg Vitamin B6, 2 mg Vitamin B12, 5.5 mg Niacin, 0.2 mg Pantothenic acid, 30 mg Betaine; \* T<sub>0</sub>: Control without the addition of date seed flour, T<sub>1</sub>: Basal feed + 2.5% date seed flour, T<sub>2</sub>: Basal feed + 5.0% date seed flour, T<sub>3</sub>: Basal feed + 7.5% date seed flour, T<sub>4</sub>: Basal feed + 10% date seed flour

## RESULTS AND DISCUSSION

The results of adding date palm flour on growth performance can be seen in Table 3. The result showed that adding date seed flour had significant effects ( $p < 0.05$ ) on final body weight, apparent digestibility of protein, and AMEn (Tables 3 and 4). In contrast, the date

seed flour had no significant effect ( $p > 0.05$ ) on the FCR and IOFC.

The feed intake results were lowest in T4, including 10% date seed flour. This result may be due to higher fiber in the date seed causing this condition. There is a difference between T<sub>0</sub> and T<sub>4</sub>, the energy and protein, where one factor caused the result of this condition.

**Table 3.** The addition of date seed flour on the growth performance of broiler chickens

Parameters	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	SEM
FI (g)	2701.88	2675.81	2719.30	2705.47	2615.67	110.06
FBW (g)	1515.96 <sup>b</sup>	1500.00 <sup>b</sup>	1530.95 <sup>b</sup>	1531.72 <sup>b</sup>	1383.29 <sup>a</sup>	28.00
FCR	1.79	1.78	1.78	1.77	1.89	0.05
IOFC (IDR/head/day)	13127.22	13037.38	13445.98	13613.48	11271.36	1073.45

FI: Feed intake, FCR: Feed conversion ratio, FBW: Final body weight, IOFC: Income over feed cost. <sup>a,b,c,d</sup> Means with different superscripts in the row differ significant ( $p < 0.05$ ). T<sub>0</sub>: Basal diet, T<sub>1</sub>: Basal diet + 2.5% date palm flour, T<sub>2</sub>: Basal diet + 5% date palm flour, T<sub>3</sub>: Basal diet + 7.5% date palm flour, T<sub>4</sub>: Basal feed + 10% date palm flour

**Table 4.** The effects of date seed flour on protein digestibility and metabolic energy of broiler chickens

Parameters	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	SEM
Apparent digestibility of protein (%)	70.20	75.06	74.48	78.29	73.31	1.82
AME (Kcal/kg)	3659.32 <sup>d</sup>	3546.80 <sup>c</sup>	3294.25 <sup>b</sup>	3490.43 <sup>c</sup>	3161.67 <sup>a</sup>	61.25
AMEn (Kcal/kg)	3046.50 <sup>d</sup>	2891.52 <sup>c</sup>	2644.06 <sup>b</sup>	2806.96 <sup>c</sup>	2521.67 <sup>a</sup>	0.02

AMEn: Apparent metabolizable energy Nitrogen corrected, AME: Apparent metabolizable energy. <sup>a,b,c,d</sup> Means with different superscripts in the row different significant ( $p < 0.05$ ). T<sub>0</sub>: Basal diet, T<sub>1</sub>: Basal diet + 2.5% date palm flour, T<sub>2</sub>: Basal diet + 5% date palm flour, T<sub>3</sub>: Basal diet + 7.5% date palm flour, T<sub>4</sub>: Basal feed + 10% date palm flour

According to Allama *et al.* (2012), if the energy and protein do not meet standard criteria, the cell tissue would convert protein into energy. Moreover, the final body weight was optimum when date seed flour was given at 7.5%, while the body weight was at its lowest in T<sub>4</sub> (10% date seed flour). Corollary, Masoudi *et al.* (2011) reported that date flour at a 10-30% level caused a reduction in final body weight. In addition, El-Faham *et al.* (2017) found that giving 10% date palm waste in feed reduced the final body weight. Accordingly, the use of date flour for more than 10% could have a significant effect on body weight. The reason can be the high crude fiber content of date seed flour. The capability of absorption is one factor on the growth performance instead feed intake and body weight (Astuti *et al.*, 2015). Furthermore, FCR is the ratio of given feed weight over animal weight gain. An increase in FCR was indicative of inefficient feed used during the experiment. In addition, the replacement of corn with 15% date pits resulted in the FCR range of 1.72-1.84 (Hammod *et al.*, 2018).

Income over feed cost can be reflected in the successful rearing management in the poultry industry. The IOFC in T<sub>3</sub> was higher than in control and other treatments. According to Muchlis *et al.* (2021), IOFC is

influenced by several factors, including feed intake, final body weight, and the selling price of broiler chickens. The FCR is key to the IOFC of broiler chickens (Prayitno *et al.*, 2019).

The apparent digestibility of protein is one of the essential indicators related to the growth performance of broiler chickens (Azizah *et al.*, 2020). The T<sub>3</sub> presented the best value for the apparent digestibility of protein. The appropriate value of the apparent digestibility indicated the high protein utilization in the broiler chickens. Furthermore, the higher result of apparent digestibility of protein can be reflected in the content of micro minerals. As El-Faki (2002) mentioned, date seeds consist of micro minerals, including iron (Fe) 7.4 mg/100 g, manganese (Mn) 2.8 mg/100 g, zinc (Zn) 1.9 mg/100 g, and copper (Cu) 1.2 mg/100 g. The Cu minerals had a main role in inhibiting microorganisms' activity in the digestive tract. Meanwhile, Zn minerals act as cofactors for protease enzymes that break down protein molecules into amino acids (Azizah *et al.*, 2020). Natsir *et al.* (2018) reported that a reduction in protein digestibility causes low body weight in broiler chickens.

Metabolic energy measured in this study is AME and AMEn. The results indicated that the higher the use of date

seed flour in the feed, the lower the AME value (Table 4). This result may be due to the high crude fiber content in date seed flour which causes broiler chickens to require more energy to digest crude fiber. Feed ingredients with high crude fiber content can reduce the digestibility value of other feed ingredients because digesting the crude fiber content requires more energy (Noviadi et al., 2012). Therefore, the nutrient contents in date seed flour affected the nutrient content in experiments.

The AMEn result ( $T_4$ : Basal feed + 10% date palm flour) was lower than that of the AME ( $T_0$ : Basal diet + 0% control) caused by nitrogen retention in AMEn. Biologically, metabolic energy is determined by nitrogen retention (Sukaryana, 2010; Table 4). In addition, Dady et al. (2015) stated that high crude fiber will cause the availability of energy in the feed to decrease. The undigested crude fiber as well as other undigested nutrients, will excrete (Siabandi et al., 2018).

## CONCLUSION

In conclusion, adding 10% date seed flour helps increase final body weight, apparent protein digestibility, and apparent metabolizable energy, without any adverse effect on the broiler chickens.

## DECLARATION

### Acknowledgments

The authors wish thanks to PT. Rumah Desa Sejahtera and Mr. Ali Baradja who have provided funding on this project. With granted number 10/01/2022.

### Author's contribution

Ilmiatus Sholichatunnisa contributed to collecting data, analysis of proximate, data analysis, and preparing the manuscript. Osfar Sjojfan contributed to the research design and supervision; Tri Eko Susilorini provided a sample and site for *in vivo* and supervision. Muhammad Halim Natsir was the supervisor. Danung Nur Adli supervised and revised the manuscript grammatically. All authors read and approved the final version of the manuscript in the present journal.

### Competing interest

No potential conflict of interest relevant to this article was reported.

### Ethical consideration

All authors have been checked the ethical issues, plagiarism, fabrication and/or falsification, double publication, and redundancy.

## REFERENCES

- Adli DN (2021a). Uses insects in poultry feed as replacement soya bean meal and fish meal in development countries: A systematic review. *Livestock Research for Rural Development*, 33: 128. Available at: <https://lrrd.cipav.org.co/lrrd33/10/33128danun.html>
- Adli DN (2021b). The effect of replacing fish meal with Sago larvae meal (SLM) on egg production and quality of laying hens. *Livestock Research for Rural Development*, 33: 94. Available at: <http://www.lrrd.org/lrrd33/7/3394danun.html>
- Allama H, Sjojfan O, Widodo E, and Prayogi HS (2012). The effect of using caterpillar flour (*Alphitobius diaperinus*) in feed on the appearance of broiler production. *Indonesian Journal of Animal Science*, 22(3): 1-8. Available at: <https://jiip.ub.ac.id/index.php/jiip/article/view/116>
- Astuti FK, Busono W, and Sjojfan O (2015). Effect of addition of liquid probiotics in feed on production appearance in broilers. *Indonesian Journal of Environment and Sustainable Development*, 6(2): 99-104. Available at: <https://jpal.ub.ac.id/index.php/jpal/article/view/192>
- Azizah TRN, Singgih DP, Setiyatwan H, Widjastuti T, and Asmara IY (2020). Increasing the utilization of ration on sentul chickens given noni fruit extract (*Morinda citrifolia*) with copper and zinc supplementation. *Journal of Tropical Animal Nutrition and Feed Science*, 2(1): 25-34. DOI: <https://www.doi.org/10.24198/jnttip.v2i1.26667>
- Bouaziz F, Ben Abdeddayem A, Koubaa M, Ellouz Ghorbel R, and Ellouz Chaabouni S (2020). Date seeds as a natural source of dietary fibres to improve texture and sensory properties of wheat bread. *Foods*, 9(6): 737. DOI: <https://www.doi.org/10.3390/foods9060737>
- BSN (2015a). Standar Nasional Indonesia (SNI) 8173.1: Broiler chickens – part 1: Starter periods. Ministry of Agriculture. Jakarta, Indonesia, pp. 23-43.
- BSN (2015b). Standar Nasional Indonesia (SNI) 8173.3: Broiler chickens – part 3: Finisher periods. Ministry of Agriculture. Jakarta, Indonesia, pp. 45-47.
- Dady Z, Mandey JS, Imbar MR, and Regar MN (2015). The value of nitrogen retention and metabolic energy in feed using fresh mulberry (*Morus alba*) leaves in broilers. *Zootec*, 36(1): 42-50. DOI: <https://www.doi.org/10.35792/zot.36.1.2016.9441>
- El-Faham AI, Ali NGM, Ali RAM, and Abdelaziz MAM (2017). Date palm waste in starter-grower diets and impact on productive and physiological performance of broilers. *Egyptian Journal of Nutrition and Feeds*, 20(2): 299-308. DOI: <https://www.doi.org/10.21608/ejnf.2017.75214>
- El-Faki MH (2002). Nutritional value of date pits for broiler. M. SC Thesis, Faculty of Animal Production, University of Khartoum, Sudan. Available at: <http://khartoumspace.uofk.edu/bitstreams/a5601d86-cc7c-4b68-9398-961b9bab2a0f/download>
- Farrell DJ (1978). Rapid determination of metabolizable energy of foods using cockerels. *British Poultry Science*, 19(1): 303-308. DOI: <https://www.doi.org/10.1080/00071667808416480>
- Giovanny L, Lestari FA, Marfira N, Ambarsari L, and Warnasih S (2019). Potency of ethanol extracts of palm seeds

- (*Phoenix dactylifera* L.) as antidiabetic with inhibition kinetics parameter. *Current Biochemistry*, 6(2): 55-67. DOI: <https://www.doi.org/10.29244/cb.6.2.1>
- Hammod AJ, Ali NA, Alkassar AM, and Jameel YJ (2018). The effect of partial replacement of maize by date pits on broiler performance. *Journal of Pure and Applied Microbiology*, 12(2): 807-813. DOI: <https://www.doi.org/10.22207/JPAM.12.2.42>
- Kresnadipayana D and Lestari D (2017). Determination of borax levels in dates (*Phoenix dactylifera*) using UV-VIS spectrophotometry method. *Wiyata Journal*, 4(1): 23-30. Available at: <http://www.ojs.iik.ac.id/index.php/wiyata/article/view/140>
- Masoudi A, Chaji M, Bojarpour M, and Mirzadeh K (2011). Effects of different levels of date pits on performance, carcass characteristics and blood parameters of broiler chickens. *Journal of Applied Animal Research*, 39(4): 399-405. DOI: <https://www.doi.org/10.1080/09712119.2011.621790>
- Muchlis A, Asmawati, Aqmal A, Hasyim Z, Reza R, Sanda E, and Resky (2021). Performance and income over feed cost (IOFC) of broiler chicken with intake of earthworm flour (*Lumbricus rubellus*) as additive in broiler basalt feed. *Journal of Integrated Animal Science and Technology*, 1(1): 7-14. Available at: <https://journal.unibos.ac.id/jitpu/article/view/1087>
- Natsir MH, Djunaidi I, Sjojfan O, Suwanto A, Puspitasari E, and Virginia LJ (2018). The effect of corn substitution with palm kernel meal treated by enzyme on production performance and carcass quality of broiler. *Buletin of Animal Science*, 42(2): 103-108. DOI: <https://www.doi.org/10.21059/buletinpeternak.v42i2.30977>
- Noviadi R, Sofiana A, and Panjaitan I (2012). The effect of using corn flour in making cassava leaf waste silage on changes in nutrition, digestibility of dry matter, crude protein and crude fiber in local rabbits. *Jurnal Penelitian Pertanian Terapan*, 12(1): 6-12. DOI: <https://www.doi.org/10.25181/jppt.v12i1.192>
- Prayitno AH, Prasetyo B, Sutirtoadi A, and Sa'roni A (2019). The effect of giving fermented tofu dregs as conventional feed on production costs of broiler ducks. *Jurnal Ilmu Peternakan Terapan*, 2(2): 50-56. Available at: <https://publikasi.poliije.ac.id/index.php/jipt/article/view/1475>
- Primurdia EG and Kusnadi J (2014). Antioxidant activity of sari kurma (*Phoenix dactylifera* L.) Probiotic drink with *L. plantarum* and *L. casei* isolates. *Jurnal Pangan dan Agroindustry*, 2(3): 98-109. Available at: <https://jpa.ub.ac.id/index.php/jpa/article/view/57>
- Regar MN and Kowel YHS (2021). Digestibility of broiler ration containing a combination of turmeric, garlic with zinc minerals. *Zootec*, 41(1): 311-316. DOI: <https://www.doi.org/10.35792/zot.41.1.2021.33862>
- Risa H, Marsudi E, and Azhar (2018). Analysis of dates plantation business feasibility (case study of barbate date gardens, aceh besar district). *Scientific Journal of Agricultural Students Unsyiah*, 3(4): 550-562. DOI: <https://www.doi.org/10.17969/jimfp.v3i4.9484>
- Setyawan MI, Sarengat W, and Bachelor TA (2019). The effect of addition of lime juice (*Citrus aurantifolia*) in drinking water on the performance of broiler chickens. *Journal of Animal Husbandry Extension Development*, 16(29): 32-38. DOI: <http://www.doi.org/10.36626/jppp.v16i29.65>
- Siabandi R, Bagau B, Imbar MR, and Regar MN (2018). Nitrogen retention and metabolic energy of broiler ration containing kepok banana peel silage flour (*Musa paradisiaca formatypica*). *Zootec*, 38(1): 226-234. DOI: <https://www.doi.org/10.35792/zot.38.1.2018.18943>
- Sjojfan O, Adli DN, Natsir MH, Nuningtyas YF, Wardani TS, Sholichatunnisa I, Ulpah SN, and Firmansyah O (2021a). Effect of dietary modified-banana-tuber meal substituting dietary corn on growth performance, carcass trait and dietary-nutrients digestibility of coloured-feather hybrid duck. *Jurnal Ilmu Ternak dan Veteriner*. 26(1):39-48. DOI: <https://www.doi.org/10.14334/jitv.v26i1.2686>
- Sjojfan O, Adli DN, Natsir MH, Nuningtyas YF, Bastomi I, and Amalia FR (2021b). The effect of increasing levels of palm kernel meal containing  $\alpha$ - $\beta$ -mannanase replacing maize to growing-finishing hybrid duck on growth performance, nutrient digestibility, carcass trait, and VFA. *Journal of the Indonesian Tropical Animal Agriculture*, 46(1): 29-39. DOI: <https://www.doi.org/10.14710/jitaa.46.1.29-39>
- Sukaryana Y (2010). Increasing metabolic energy of fermentation products mixed palm kernel meal and rice bran. *Jurnal Penelitian Pertanian Terapan*, 10(2): 138-143. DOI: <https://www.doi.org/10.25181/jppt.v10i2.256>
- Tareen MH, Wagan R, Siyal FA, Babazadeh D, Bhutto ZA, Arain MA, and Saeed M (2017). Effect of various levels of date palm kernel on growth performance of broilers. *Veterinary World*, 10(2): 227-232. DOI: <https://www.doi.org/10.14202/vetworld.2017.227-232>