










# Efficiency of Recycled Plastic Bedding Material and Gender in Improvement of Productive Traits, Physiological, and Immunological Parameters of Hybrid Broiler Chickens

Mohamed A. EL-Masry<sup>1</sup> , Magdy S. Hassan<sup>2</sup> , AbdelSatar A. Arafa<sup>3\*</sup> , Tarek M. El-Afifi<sup>4</sup> , Ahmed M. Bealish<sup>2</sup> ,  
Magdy M. Ouda<sup>2</sup> , Ibrahim A. Fathey<sup>2</sup> , Hanan A. Fahmy<sup>3</sup>  and Hanaa K. Abd El-Atty<sup>2</sup> 

<sup>1</sup>Misr Organic Food Industry Company, Dokki, Giza, Egypt; <sup>2</sup>Animal Production Research Institute (APRI), Agriculture Research Center (ARC), Dokki, Giza, Egypt; <sup>3</sup>Animal Health Research Institute (AHRI), Agriculture Research Center (ARC), Dokki, Giza, Egypt; <sup>4</sup>Regional Center for Food and Feed, Agriculture Research Center (ARC) Giza, Egypt.

\*Corresponding author's E-mails: [abd.arafa@hotmail.com](mailto:abd.arafa@hotmail.com); [abd.arafa@gmail.com](mailto:abd.arafa@gmail.com), [abd.arafa@ahri.gov.eg](mailto:abd.arafa@ahri.gov.eg)

Received: March 31, 2024, Revised: April 23, 2024, Accepted: May 15, 2024, Published: June 30, 2024

## ABSTRACT

Litter management is important for poultry housing husbandry and affects chicken performance. The present study evaluated the effect of bedding material and gender on the productive, physiological, and immunological performance of a new hybrid chicken (WINZY Line 105) under cold stress for 56 days of age. A total of 540 one-day-old broiler chicks were divided into two groups. The groups, including 270 males (M) and 270 females (F) were further divided into two sub-groups, including sawdust litter (SL) and plastic slatted floor (PSF), three replicates, and 45 chickens each. Broiler chickens were raised during the winter with an average temperature of 10°C for 56 days. Productive, physiological, and immunological performance parameters were measured. The obtained results indicated that M reared on PSF (M x PSF) recorded the highest values in body weight (BW), body weight gain (BWG), carcass characteristics, and the best values in feed conversion compared with other interaction groups during all experimental periods. In addition, F reared on PSF had higher BW, carcass, and thigh percentages, low feed intake (FI), and best feed conversion compared to females reared on SL. The M reared on PSF had the highest hepatic enzymes except AST which was higher in F reared on PSF than other treatments. However, renal function biomarkers (Creatine, Uric acid, Urea) were higher in both M and F that were reared on SL than those reared on PSF. Moreover, there was a significant interaction detected for antibody titers against avian influenza (H5) and Newcastle disease at 21 days of age suggesting that the highest values observed for M reared on PSF (M x PSF) compared with other interaction groups during the experimental period, and it was higher in F that reared on PSF than F reared on SL at 21 days of age. It can be concluded that plastic slatted floors could be an alternative to substitute wood shavings to raise broiler chickens since it was efficient from the perspective of environmental conditions and production rates.

**Keywords:** Bedding material, Environmental adaptation, Hybrid chicken, Immunological parameter, Productive trait, Plastic slatted floor, Performance

## INTRODUCTION

Bedding management is critical for animal welfare, especially in intensive production systems. Litter protects the chickens from the rigid ground, and facilitates the evaporation of urine, excreta, and spilled water. To improve the health and welfare of chickens, they should be kept on a dry floor. A variety of factors are used in broiler production to meet these goals. The main factors are the type of litter material used, housing system, litter control methods, and bird water balance (Collett, 2012).

The purpose of bedding material included excreta and moisture absorption, aerobic decomposition of excrement, and heat insulation (Shepherd and Fairchild, 2010). Additionally, it influences the body weight (BW) and immunity of broiler chicks, which reflects on growth rate and carcass quality like carcass body weight percent. Broilers reared on sand and paper roll floors perform similarly to those reared on wood shavings (Bilgili et al., 2009; Toghyani et al., 2010).

Broiler performance can be enhanced through practical litter management to ensure a healthy atmosphere

in their houses by controlling ammonia concentrations, and increasing immunological responses against various diseases (Beker et al., 2004; Miles et al., 2004; De Jong et al., 2014; Wei et al., 2015). In addition, heat exhaustion puts broiler breeding in danger (Liverpool-Tasie et al., 2019).

The most common materials used as litter in commercial broiler production are wood shavings and sawdust; recently, plastic floors have been effective in providing a healthy environment and higher production rates. Wood shavings and plastic floors demonstrated better performance with males than females at 42 days of age including weight increase, feed intake, and feed conversion (Almeida et al. 2017).

Climate change negatively impacted both direct and indirect agricultural production systems and people's food security, especially in dry areas like Sub-Saharan Africa (Thompson et al., 2010). The Food and Agricultural Organization (FAO, 2011) advised increasing public awareness of how climate change affects food security and nutrition, the mechanisms affecting food security, and how to adapt to climate change.

This study aims to investigate the effect of two different bedding materials (sawdust litter and plastic slatted floors) on the growth performance and immunological parameters of both males and females of the new hybrid chickens WINZY Line 105.

## MATERIALS AND METHODS

### Ethical approval

The animal study protocol was conducted with permission and approved by the Review Board of Animal Production Research Institute (APRI), Agriculture Research Center, Ministry of Agriculture, Dokki, Giza, Egypt. All experiments were conducted according to the relevant guidelines and regulations of the Ethical Committee coded: 202110

### Broiler chickens

A new hybrid breed broiler that originated from a crossing between the local native Egyptian breed (Fayomy) PP line and the French line breed (Sasso) GG line. The average body weight for males is 2.2 kg, while for females it was 1.750 kg at 52 days of age. This line was characterized by a sex phenotype difference where the color of females was reddish brown and can be easily differentiated from the yellowish-grey color of males at one day old.

### Experimental design

The current study was conducted in collaboration with Misr Organic Food Industry Company, Animal Production Research Institute Animal Health Research Institute, Agricultural Research Center, Ministry of Agriculture, Egypt. This study was conducted at the facility of poultry breeding station, Anshas, Sharkaia, Egypt. The chicks were hatched from the fifth generation of cross-type chickens (WINZY line 105) of both sexes. A total of 540 chicks were reared from one day old up to 56 days old. The chicks were divided into two groups of 270 males (M) and 270 females (F). Groups were further divided into two sub-groups based on the type of litter: sawdust litter (SL) and plastic slatted floor (PSF), with 3 replicates of 45 chickens each under the same management conditions including ventilation system, lighting program, temperature inside houses, feeding formulae, and vaccination programs. The composition and calculated analysis of the basic diet was done according to the NRC (1994) and presented in Table 1. The chickens, house facility, and plastic-slatted floor were supplied by Misr Organic Food Industry Company.

### Floor design

The floors were distributed in rooms at the same height in each room. Broiler chickens were reared on two types of bedding material (sawdust litter and plastic slatted floor). Sawdust litter was distributed with height (7 cm) and considered as control groups. The plastic slatted floor was designed with narrow holes measured 2-3 millimeters thick. Water and feed diet were provided *ad libitum*. Broiler chickens in this experiment were reared during the winter season (December 2021 and January 2022) under extreme cold conditions. The ambient temperature in the daytime reached 10°C and fell to 2°C at night. The chicks were reared in a semi-closed, controlled house with a nearly fixed temperature inside to maintain chickens during growth.

### Effective operation

Live body weight (BW) and feed intake (FI) were recorded weekly, consequently body weight gain (BWG) and the feed conversion (FC) ratio were calculated too. Daily mortalities were counted, and a weekly mortality rate was computed.

### Features of slaughter and carcass

After eight weeks, chickens were kept fasting for 5-6 hours to keep the corps of the chickens empty at slaughtering time, then three chickens from each treatment were slaughtered by slitting the jugular vein, de-feathered,

eviscerated, weighted, and the percentages of live body weights were calculated. The breast, thigh, and giblets

were excised, weighed, and expressed as percentages to live body weights.

**Table 1.** Composition of basal diet of the hybrid broiler (WINZY Line 105) affected by gender and plastic-based bedding material from day one to eight weeks of age

Ingredients (%)	Starter (1-21 days)	Grower (22-42 days)	Finisher (43-56 days)
Yellow corn	56.00	61.65	65.25
Soya bean meal 44%	32.50	25.00	21.00
Corn gluten meal 62%	6.00	7.00	7.00
Cotton seed oil	2.00	2.50	3.25
Di-calcium phosphate	0.80	1.00	1.00
Limestone	1.85	2.00	1.75
Salt	0.25	0.25	0.25
Mineral and vitamin mix*	0.30	0.30	0.30
DL-Methionine	0.15	0.15	0.10
L-lysine	0.15	0.15	0.10
Total	100	100	100
<b>Calculated analysis</b>			
Crude protein (%)	23.08	20.88	19.29
ME (kcal/kg diet)	3004.95	3104.62	3201.57
Crude fiber (%)	3.61	3.24	3.08
Calcium (%)	1.00	1.00	0.97
Available phosphorus (%)	0.55	0.56	0.54
Methionine (%)	0.53	0.52	0.46
Lysine (%)	1.23	1.07	0.95
Methionine+Cysteine (%)	0.91	0.87	0.79

\*Vitamin and mineral mix. mix: each 3kg contains: Vit. A, 12000000 IU; Vit. D3, 2000000 IU; Vit. E, 10 g; Vit. K, 2.0 g; Vit. B1, 1g; Vit. B2, 5g; Vit. B6, 1.5 g; Vit. B12, 10 mg; Folic acid, 1g; Biotin, 50mg; Pantothenic acid, 10 g; Nicotinic acid, 30 g; Choline chloride, 250 g; Mn, 60g; Fe 30, g; Zn, 50 g; Cu, 10g; I, 1 g; Co 100 mg; Se, 100 mg; Anti-oxidant, 10 g, and complete to 3.0 kg by Calcium Carbonate. The data in the table was calculated according to NRC (1994).

### Blood parameters

Blood parameters were measured at the Reference Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Giza, Egypt. Blood testing was carried out to evaluate physiological performance in the tested groups. Blood samples were taken from 18 chickens/treatment after slaughtering at 56 days of age from each group in collecting tubes containing heparin as an anticoagulant to obtain whole blood as 1 mL per individual bird. Then, it was centrifuged for 15 min at 3000 rpm and stored at -20°C. Blood testing parameters conducted in this study included: liver function testing the concentrations of two enzymes, aspartate transaminase (AST) and alanine transaminase (ALT), and measuring glucose level, alkaline phosphatase (ALP), and kidney function testing creatinine (CRE), uric acid and urea by calorimetric methods using commercial kits of Bio Diagnostic Co., Egypt, following the manufacturer's instructions.

### Immunological parameters

On days 10 and 21, blood samples were collected to determine immunological parameters including antibody titers against avian influenza virus (H5) and Newcastle disease (ND). Blood samples were taken on days 10 and 21 from 18 chickens from each group using sterile plastic syringes to collect blood from the wing vein as 1 mL per individual bird then left for complete agglutination to separate serum. The collected sera were transferred to Eppendorf tubes and centrifuged at 3000 rpm to separate clear sera. Serum samples were tested to determine the immunological response after vaccination by measuring antibody titers against avian influenza virus (AIV-H5) and Newcastle disease virus (NDV) using Hemagglutination inhibition test (HI) according to standard protocol (WOAH, 2021).

### Statistical analysis

The experiment data were statistically examined by analysis of variance according to Mead (2002) using ANOVA procedures of SAS (SAS, 2011). The statistical model was used as the following formula.

$$Y_{ij} = \mu + Z_i + S_j + (ZS)_{ij} + e_{ij}$$

Where  $Y_{ij}$  is an observation,  $\mu$  is the overall mean,  $Z_i$  is the effect of the sex groups, ( $i$  is equal to 1 and 2),  $S_j$  is the effect of bedding materials ( $j$  is equal to 1 and 2),  $(ZS)_{ij}$  is the interaction effect between sex and bedding materials ( $ij = 1, 2, \dots, 4$ ),  $e_{ij}$  refer to random error.

The differences between means were tested by using Duncan's multiple-range test procedures (Duncan, 1955), and  $p$ -values less than 0.05 were considered significant. The percentage values were subjected to sine transformation before analyzing variance. Means were present after recalculating from the transformed value to percentages.

## RESULTS

### Body weight and body weight gain

The effects of sex and bedding material on BW and BWG are illustrated in Table 2. During all the experimental periods, males (M) had significantly higher BW and BWG values than females (F). However, the bedding material has a low impact on the BW. Chickens raised on plastic slatted floors (PSF) had higher BWGs at different ages (1-8 weeks) than chickens kept on sawdust litter (SL). The results indicated that a significantly higher interaction was found for BW and BWG, indicating that the highest values were observed for M reared on PSF compared with those of the other interaction groups during all experimental periods and this interaction was higher for F reared on PSF than for F reared on SL ( $p < 0.05$ ).

### Feed intake and feed conversion

The effect of sex and bedding type on FI and FC are shown in Table 3. Significant increases in FI and FC among males and females were found during all the experimental periods ( $p < 0.05$ ). Compared with females, males presented a higher FI during all experimental periods and improved FC by 0.2% during the whole 8 weeks and 0.25% during (4-8 weeks). No significant differences were found in FI during all experimental periods for chickens reared on either floor type (SL or PSF), while chickens reared on PSF had higher FCs by 0.08% and 0.06%, respectively, than chickens housed on SL at 4-8 and 1-8 weeks of age ( $p < 0.05$ ). The interaction effects demonstrated that the FI of males raised on both types of floors did not differ significantly. However, compared to females raised on SL, those raised on PSF had lower FI and higher FC. For all the experimental periods, the FC values for males raised on PSF were

higher than those for the other interaction groups ( $p < 0.05$ ).

### Carcass traits of several internal organs

As shown in Table 4, males had significantly higher live body weight (LBW) than females including carcass 2.3%, breast 0.97%, and thigh 0.8%, while giblet percentages were not significantly different ( $p < 0.05$ ). Compared with those raised on sawdust litter, the chickens raised on plastic slatted floors had a substantially higher percentage of carcass by 1.55%, breast by 0.83%, and thigh by 0.98% ( $p < 0.05$ ). The interaction effects indicated no significant differences in LBW between males reared on both floor types (SL and PSF;  $p < 0.05$ ). However, females reared on PSF had higher LBW (carcass by 1.2% and thigh by 1.18%) than females reared on SL. Among the males reared on SL, those reared on PSF had the highest LBW (carcass by 1.9%, breast by 0.84% and thigh by 0.79%;  $p < 0.05$ ). Furthermore, there was no significant difference in giblet percentages among the groups ( $p > 0.05$ ).

### Blood parameters

The hepatic and renal function biomarkers affected by the sex and rearing system are presented in Table 5. Alanine transaminase (ALT) and glucose levels were significantly higher in males than in females. However, renal function biomarkers (CRE, uric acid, and urea) were significantly higher in F than in M ( $p < 0.05$ ). Chickens that were reared on PSF had the highest hepatic function biomarkers AST and ALT and the lowest renal function biomarkers ( $p < 0.05$ ). The interaction effects indicated also that M reared on PSFs had the highest hepatic function biomarkers, except for AST, which was higher in F reared on PSFs than in those of other groups ( $p < 0.05$ ). However, renal function biomarkers were higher in both M and F chicks reared on SL.

### Immunological performance

According to the data in Figure 1, M had significantly higher antibody titers against avian influenza AI (H5) and ND at the 10<sup>th</sup> and 21<sup>st</sup> days of age than females ( $p < 0.05$ ). Moreover, chickens reared on PSF had higher antibody titers against AI (H5) and ND on the 21<sup>st</sup> day of age than those reared on SL. The data revealed that there was a higher interaction effect on the antibody titers against AI (H5) and ND at 10<sup>th</sup> and 21<sup>st</sup> days of age, suggesting that the highest values were observed for M reared on PSF compared with those of the other interaction groups during the whole experimental period ( $p < 0.05$ ).

**Table 2.** Body weight and body weight gain of the hybrid broiler (WINZY Line 105) affected by gender and plastic-based bedding material from day one to eight weeks of age

Items		Body weight			Weight gain		
		1d	4 W	8 W	1d-4 W	4-8 W	1d-8 W
Gender	M	33.64±0.68	594.48±15.52 <sup>a</sup>	1813.78±22.04 <sup>a</sup>	553.36±10.30 <sup>a</sup>	1234.61±22.70 <sup>a</sup>	1787.97±23.88 <sup>a</sup>
	F	33.67±0.26	510.30±10.01 <sup>b</sup>	1508.50±21.81 <sup>b</sup>	476.63±9.89 <sup>b</sup>	998.20±16.30 <sup>b</sup>	1474.83±22.12 <sup>b</sup>
	P Value	0.956	0.006	0.001	0.001	0.001	0.001
Bedding materials	SL	33.66±0.48	544.00±45.27	1646.33±166.40	510.34±45.23	1102.33±122.16 <sup>b</sup>	1612.67±166.40 <sup>b</sup>
	PSF	33.65±0.55	560.78±46.03	1683.78±171.31	519.65±38.38	1130.48±130.75 <sup>a</sup>	1650.13±171.31 <sup>a</sup>
	P Value	0.983	0.609	0.055	0.590	0.028	0.041
<b>Interaction between gender and bedding materials</b>							
Gender	Bedding materials						
M	SL	33.64±0.76	585.44±10.52 <sup>a</sup>	1800.67±7.40 <sup>b</sup>	551.80±3.54 <sup>a</sup>	1215.22±11.08 <sup>b</sup>	1767.02±8.70 <sup>b</sup>
	PSF	33.64±0.76	603.51±1.33 <sup>a</sup>	1842.56±6.38 <sup>a</sup>	554.91±9.41 <sup>a</sup>	1254.00±10.22 <sup>a</sup>	1808.91±6.43 <sup>a</sup>
F	SL	33.68±0.07	502.56±6.57 <sup>c</sup>	1492.00±10.00 <sup>d</sup>	468.87±11.12 <sup>b</sup>	989.44±6.59 <sup>c</sup>	1458.32±10.93 <sup>d</sup>
	PSF	33.66±0.41	518.04±5.12 <sup>bc</sup>	1525.00±9.95 <sup>c</sup>	484.38±10.93 <sup>b</sup>	1006.9±5.06 <sup>c</sup>	1491.34±11.68 <sup>c</sup>
P Value.		0.973	0.022	0.007	0.002	0.003	0.002

<sup>a, b, c, d</sup> Means bearing different superscript letters within the same column were significantly different (p < 0.05). d: Day, W: week, M: Males, F: Females, SL: Sawdust litter, PSF: Plastic slatted floor.

**Table 3.** Feed intake and feed conversion of the hybrid broiler chickens affected by gender and plastic-based bedding material from day one to eight weeks of age

Items		Body weight			Weight gain		
		1 d – 4 wk.	4 wk.- 8 wk.	1d- 8wk.	1d-4 W	4-8 W	1d -8 W
Gender	M	1253.85±11.37 <sup>a</sup>	2799.30±43.03 <sup>a</sup>	4053.15±51.94 <sup>a</sup>	2.27±0.05 <sup>b</sup>	2.27±0.04 <sup>b</sup>	2.27±0.04 <sup>b</sup>
	F	1137.58±20.33 <sup>b</sup>	2511.04±50.61 <sup>b</sup>	3648.62±54.25 <sup>b</sup>	2.39±0.06 <sup>a</sup>	2.52±0.08 <sup>a</sup>	2.47±0.07 <sup>a</sup>
	P Value	0.023	0.007	0.005	0.032	0.008	0.006
Bedding materials	SL	1189.67±66.67	2666.94±130.0	3856.62±199.52	2.34±0.08	2.43±0.10 <sup>a</sup>	2.40±0.12 <sup>a</sup>
	PSF	1201.76±60.33	2643.40±183.22	3845.16±222.11	2.32±0.09	2.35±0.12 <sup>b</sup>	2.34±0.09 <sup>b</sup>
	P Value	0.678	0.057	0.122	0.321	0.046	0.033
<b>Interaction between gender and bedding materials</b>							
Gender	Bedding materials						
M	SL	1253.09±5.12 <sup>a</sup>	2784.69±30.3 <sup>a</sup>	4037.79±28.66 <sup>a</sup>	2.27±0.04 <sup>c</sup>	2.29±0.02 <sup>c</sup>	2.29±0.03 <sup>c</sup>
	PSF	1254.61±11.18 <sup>a</sup>	2813.91±45.39 <sup>a</sup>	4068.52±62.22 <sup>a</sup>	2.26±0.05 <sup>c</sup>	2.24±0.04 <sup>d</sup>	2.25±0.05 <sup>d</sup>
F	SL	1126.25±6.51 <sup>b</sup>	2549.19±14.55 <sup>b</sup>	3675.44±17.17 <sup>b</sup>	2.40±0.05 <sup>a</sup>	2.58±0.03 <sup>a</sup>	2.52±0.02 <sup>a</sup>
	PSF	1148.91±14.61 <sup>b</sup>	2472.89±25.62 <sup>c</sup>	3621.80±66.54 <sup>c</sup>	2.37±0.06 <sup>b</sup>	2.46±0.05 <sup>b</sup>	2.43±0.05 <sup>b</sup>
P Value.		0.037	0.008	0.009	0.042	0.009	0.008

<sup>a, b, c, d</sup> Means bearing different superscript letters within the same column were significantly different (p < 0.05). d: Day, W: Week, M: Males, F: Females, SL: Sawdust litter, PSF: Plastic slatted floor

**Table 4.** Carcass traits of the hybrid broiler (WINZY Line 105) affected by gender and plastic-based bedding materials from day one to eight weeks of age


Items		Body weight (gm)	Carcass (%)	Breast (%)	Thigh (%)	Giblets (%)
Gender	M	1901.67±12.69 <sup>a</sup>	78.25±1.95 <sup>a</sup>	25.45±1.08 <sup>a</sup>	28.02±0.54 <sup>a</sup>	4.86±0.09
	F	1771.67±11.19 <sup>b</sup>	76.28±1.88 <sup>b</sup>	24.48±0.64 <sup>b</sup>	27.22±1.12 <sup>b</sup>	4.78±0.15
	P Value	0.006	0.033	0.022	0.036	0.151
Bedding materials	SL	1820±14.62	76.49±2.23 <sup>b</sup>	24.55±0.78 <sup>b</sup>	27.13±1.09 <sup>b</sup>	4.80±0.16
	PSF	1853.33±54.04	78.04±1.51 <sup>a</sup>	25.38±0.64 <sup>a</sup>	28.11±0.48 <sup>a</sup>	4.84±0.09
	P Value	0.336	0.047	0.041	0.040	0.611
Effect of interaction between gender and bedding materials						
Gender	Bedding materials					
M	SL	1898.33±30.12 <sup>a</sup>	77.31±1.43 <sup>bc</sup>	25.03±1.02 <sup>bc</sup>	27.62±0.12 <sup>b</sup>	4.83±0.10
	PSL	1905.00±32.23 <sup>a</sup>	79.19±0.99 <sup>a</sup>	25.87±0.88 <sup>a</sup>	28.41±0.46 <sup>a</sup>	4.88±0.09
F	SL	1741.67±12.52 <sup>c</sup>	75.67±2.85 <sup>d</sup>	24.07±0.56 <sup>d</sup>	26.63±1.23 <sup>c</sup>	4.76±0.22
	PSL	1801.67±14.46 <sup>b</sup>	76.89±0.88 <sup>cd</sup>	24.90±1.12 <sup>c</sup>	27.81±0.45 <sup>b</sup>	4.79±0.08
P Value.		0.024	0.031	0.042	0.032	0.345

<sup>a, b, c, d</sup> Means bearing different superscript letters within the same column were significantly different (p < 0.05). d: Day, W:week, M: Males, F: Females, SL: Sawdust litter, PSF: Plastic slatted floor

**Table 5.** Blood parameters of the hybrid broiler (WINZY Line 105) as affected by gender and plastic-based bedding material from day one to eight weeks of age

Items		AST (U/L)	ALT (U/L)	Glucose (mg/100ml)	ALK (U/L)	CRE (mg/dl)	Uric acid (mg/dl)	Urea (mg/dl)
Gender	M	67.10±4.19	59.55±2.02 <sup>a</sup>	81.05±3.54 <sup>a</sup>	110.44±4.94	0.81±0.08 <sup>b</sup>	3.80±.62 <sup>b</sup>	4.81±1.16 <sup>b</sup>
	F	65.70±4.33	56.38±2.17 <sup>b</sup>	78.14±1.40 <sup>b</sup>	109.37±4.2	0.89±0.11 <sup>a</sup>	4.23±.49 <sup>a</sup>	5.11±0.86 <sup>a</sup>
	P Value	0.422	0.007	0.004	0.654	0.002	0.009	0.005
Bedding materials	SL	64.07±3.76 <sup>b</sup>	56.92±2.16 <sup>b</sup>	78.07±2.86 <sup>b</sup>	105.78±0.88 <sup>b</sup>	0.93±0.06 <sup>a</sup>	4.31±.60 <sup>a</sup>	5.29±0.71 <sup>a</sup>
	PSF	68.73±3.72 <sup>a</sup>	59.02±2.51 <sup>a</sup>	81.11±2.61 <sup>a</sup>	114.03±1.26 <sup>a</sup>	0.77±0.18 <sup>b</sup>	3.72±.42 <sup>b</sup>	4.63±1.11 <sup>b</sup>
	P Value	0.003	0.023	0.033	0.005	0.004	0.003	0.003
Effect of interaction between gender and bedding materials								
Gender	Bedding materials							
M	SL	66.47±3.31 <sup>b</sup>	58.77±1.00 <sup>b</sup>	78.76±2.83 <sup>bc</sup>	105.98±0.65 <sup>b</sup>	0.88±0.03 <sup>b</sup>	4.48±.68 <sup>a</sup>	5.23±0.68 <sup>a</sup>
	PSF	67.73±3.5 <sup>b</sup>	60.33±1.71 <sup>a</sup>	83.33±1.38 <sup>a</sup>	114.90±0.66 <sup>a</sup>	0.73±0.06 <sup>d</sup>	3.98±0.39 <sup>b</sup>	4.39±.54 <sup>c</sup>
F	SL	61.67±1.70 <sup>c</sup>	55.07±1.28 <sup>c</sup>	77.38±1.19 <sup>c</sup>	105.57±1.04 <sup>b</sup>	0.98±0.03 <sup>a</sup>	4.14±0.41 <sup>ab</sup>	5.34±.88 <sup>a</sup>
	PSF	69.73±3.31 <sup>a</sup>	57.70±2.28 <sup>bc</sup>	78.89±1.14 <sup>b</sup>	113.16±1.05 <sup>a</sup>	0.80±0.02 <sup>c</sup>	3.46±0.28 <sup>c</sup>	4.88±.74 <sup>b</sup>
P Value.		0.023	0.016	0.008	0.007	0.006	0.005	0.006

<sup>a, b, c, d</sup> Means bearing different superscript letters within the same column were significantly different (p < 0.05). d: Day, W: Week, M: Males, F: Females, SL: Sawdust litter, PSF: Plastic slatted floor



**Figure 1.** Antibody titers against avian influenza and Newcastle disease at days 10<sup>th</sup> and 21<sup>st</sup> of the age in hybrid broiler chickens (WINZY Line 105) as affected by gender and plastic-based bedding materials from day one to eight weeks of age. M (SL): Male (Sawdust Litter). M (PSF): Male (Plastic Slatted Floor). F (SL): Female Sawdust Litter. F (PSF): Female (Plastic Slatted Floor). \*:  $p < 0.05$ .

## DISCUSSION

Litter or bedding material is an important constituent of the poultry industry and growth performance, it primarily affects growth parameters such as BW and BWG (Almeida *et al.* 2017). In this study, the chickens reared on the plastic-slatted floor had better growth performance than those in the wood shavings. The perforated plastic floors were considered to be a good alternative to wood shavings when rearing broiler chickens both sex male and female (Almeida *et al.* 2018). They were effective at promoting a better-quality environment and superior production rates.

Females and males grown on plastic floor had a BWG significantly higher than that reported in wood shavings. Similar results were obtained in a previous study; when comparing groups reared on different floor materials, the growth of chickens reared on a plastic-slatted floor was higher in terms of final BW, BWG, and growth rate than other flooring systems (Almeida *et al.*, 2018; Çavuşoğlu *et al.* 2018).

Feed intake in females of Winzy line 105 was recorded as 3621.8g and 3675.4g at 8 weeks of age for rearing on PSF and SL respectively, while for males was 4068.5g and 4037.8g respectively. In a previous study, males grown on plastic floors were heavier than males raised on wood shavings, and females raised on plastic

floors were also heavier than those reared on the wood shavings floor.

Manning *et al.* (2007) indicated the level of water usage may be influenced by the litter quality and the degree of absorbency of the litter material.

The higher final BW and total FI of broilers in the groups raised in fully or partially slatted flooring designs were significantly higher than those of broiler chickens in the deep litter flooring system (Topal and Petek, 2021).

Carcass body weight for females reared on PSF was higher than SL in both sexes. That was agreed with Kralik *et al.* (2015), who indicated that sex had a higher significant impact on the live weight, carcass weight, and weight of the main body parts. It is also preferable for individuals of younger ages, to be reared on a plastic-slatted flooring system (Passini *et al.*, 2012). Almeida *et al.* (2017) verified that females raised on plastic flooring had a heavier carcass weight than those raised on wood shavings, but male broiler chickens had a heavier carcass weight than females. Slat flooring systems showed preference in younger ages at slaughter (Çavuşoğlu *et al.*, 2018). Poultry reared on plastic floor had higher live weights and carcass weights for males than females (Almeida *et al.*, 2018).

Abo Ghanima *et al.* (2020) investigated the effects of three litter-rearing systems including wood-shaving litter,

perforated plastic slatted-rearing systems, and cage-rearing systems, they reported that chickens raised on wood-shaving litter had lower dressing percentages than chickens raised on plastic-slatted floors. They reported that the style of flooring had no significant impact on the relative weights of the liver and heart. Broilers raised on different types of floors indicated nonsignificant differences in LBW and giblets, according to Farghly et al. (2021a; 2021b).

Compared to chickens reared on sawdust, chickens reared on perforated plastic floors had significantly heavier carcasses. According to Al-Nasseri et al. (2021), an increase in the live body weight of chickens reared on plastic floors due to general improvement in environmental conditions, including improved air quality and heat relief due to decreased dust content and decreased moisture in the floor, leading to increased performance and growth parameters. The plastic-slatted floor system allows the chickens' feet to be in direct contact with the ground, this allows high heat conductivity in the environment and facilitates airflow around the chickens. Blood parameters like AST and ALT were higher in both sexes reared on PSF than SL. These results support those of Wang et al. (2015), who reported that chickens reared in litter-based systems had lower apparent ileal digestible energy intake than those reared on the Net rearing system during the first 3 weeks of age.

The results also indicated that monitoring blood parameters was important for assessing stress and immune response to stressors in poultry (Saeed et al., 2019; Nwaigwe et al., 2020).

The increased significant interaction effect on the antibody titers against AI (H5) and ND on days 10 and 21 of age was observed in this study. Results suggested that the highest values were observed for M reared on PSF compared with those of the other interaction groups during the whole experimental period. There were higher antibody titers in female individuals reared on PSF than in female individuals reared on SL on the 21<sup>st</sup> day of age which indicated the desirable effect of PSF on immunity than the SL.

The level of AIV-H5 and NDV antibodies in chickens reared on PSF compared with those reared on sawdust litter SL revealed that antibody titers against NDV and AI were elevated in the PSF groups. These results indicated a significant increase in the level of antibodies against both viruses and increased their vaccination response and ability for stress control. Previous results suggested that rearing chickens in cages and on plastic-slatted floors could enhance immunity (Sogunle et al., 2008).

## CONCLUSION

The plastic slatted floors were effective in terms of environmental conditions and production rates, and they are suitable replacements for wood sawdust during the growing period of broilers chickens. This flooring system would enhance production rates and immunity under adverse climatic conditions in chickens. The production efficiency of the hybrid breed used in this study was enhanced using a plastic slatted floor compared to the traditional sawdust floor, especially for body weight gain and antibody titers against ND and AI.

## DECLARATIONS

### Ethical considerations

The article was written originally by authors from the obtained original data and it was not submitted or published totally or even partially in other publications. The text article is checked by a well-known plagiarism checker software before submission to the journal.

### Acknowledgments

The authors are grateful to Misr Organic Food Industry Company for its funding support for this study.

### Authors' contributions

Mohamed EL-masry (ME) and Magdy Hassan (MH) designed, performed, and followed up the experiment. Ahmed Bealish (AB), Magdy Ouda (MO), and Ibrahim Fathey (IF) ran the experiment and conceived the study. Tarek El-Afifi (TE) and Abdelsatar Arafa (AA) did the laboratory work. Hanaa Abd El-Atty (HA) wrote the original draft. Hanaa Abd El-Atty (HA), Abdelsatar Arafa (AA), and Hanan Fahmy (HF) participated in the manuscript draft review, coordination, and editing. All authors read and approved the final manuscript.

### Competing interests

The authors declare that they have no conflict of interest.

### Funding

Misr Organic Food Industry Company, Egypt funded this study by providing the hybrid chicken of the 5<sup>th</sup> generation (WINZY Line 105) and germinated feed (barley + sorghum).

### Availability of data and materials



The data that support the findings of this study are available from the corresponding author upon reasonable request.

## REFERENCES

- Abo Ghanima MM, Abd El-Hack ME, Othman SI, Taha AE, Allam AA, and Eid Abdel-Moneim AM (2020). Impact of different rearing systems on growth, carcass traits, oxidative stress biomarkers, and humoral immunity of broilers exposed to heat stress. *Poultry Science*, 99(6): 3070-3078. DOI: <https://www.doi.org/10.1016/j.psj.2020.03.011>
- Almeida EA, De Souza LFA, Sant'Anna AC, Bahiense RN, Macari M, and Furlan RL (2017). Poultry rearing on perforated plastic floors and the effect on air quality, growth performance, and carcass injuries-experiment 1: Thermal comfort. *Poultry Science*, 96(9): 3155-3162. DOI: <https://www.doi.org/10.3382/ps/pex131>
- Almeida EA, Sant'Anna AC, Crowe T, Macari M, and Furlan RL (2018). Poultry rearing on perforated plastic floors and the effect on air quality, growth performance, and carcass injuries- Experiment 2: Heat stress situation. *Poultry Science*, 97(6): 1954-1960. DOI: <http://www.doi.org/10.3382/ps/pey048>
- Al-Nasser ANI, Taha AT, and Hasan ATK (2021). Effects of different broiler flooring systems on surface temperature, air quality and carcass characters of broilers. *IOP Conference Series of Earth and Environmental Science*, 735: 012011. Available at: <https://ui.adsabs.harvard.edu/abs/2021E%26ES..735a2011A/abstract>
- Beker A, Vanhooser SL, Swartzlander JH, and Teeter RG (2004). Atmospheric ammonia concentration effects on broiler growth and performance. *Journal of Applied Poultry Research*, 13(1): 5-9. DOI: <https://www.doi.org/10.1093/japr/13.1.5>
- Bilgili SF, Hess JB, Blake JP, Macklin KS, Saenmahayak B, and Sibley JL (2009). Influence of bedding material on footpad dermatitis in broiler chickens. *The Journal of Applied Poultry Research*, 18(3): 583-589. DOI: <https://www.doi.org/10.3382/JAPR.2009-00023>
- Çavuşoğlu E, Petek M, Abdourhamane İM, Akkoc A, and Topal E. Effects of different floor housing systems on the welfare of fast-growing broilers with an extended fattening period. *Archives Animal Breeding*, 61(1): 9-16. DOI: [https://www.doi.org/10.5194/aab-61-9-2018\\_2018](https://www.doi.org/10.5194/aab-61-9-2018_2018)
- Collett SR (2012). Nutrition and wet litter problems in poultry. *Animal Feed Science and Technology*, 173(1-2): 65-75. DOI: <https://www.doi.org/10.1016/j.anifeedsci.2011.12.013>
- De Jong IC, Gunnink H, and van Harn J (2014). Wet litter not only induces footpad dermatitis but also reduces overall welfare, technical performance and carcass yield in broiler chickens. *Journal of Applied Poultry Research*, 23(1): 51-58. DOI: <https://www.doi.org/10.3382/japr.2013-00803>
- Duncan DB (1955). Multiple range and multiple F tests. *Biometrics*, 11(1): 141. DOI: <https://www.doi.org/10.2307/3001478>
- Food and agriculture organization of the United Nations (FAO) (2011). Climate change mitigation finance for smallholder agriculture, pp. 1-88. Available at: <https://www.fao.org/climatechange/29763-0daebae838c70f713da780982f16e8d9.pdf>
- Farghly MFA, Mahrose KM, Cooper RG, Metwally KA, Abougabal MS, and El-Ratel IT (2021a). Use of available crop by-products as alternative bedding materials to wheat straw for rearing broilers. *Animal: An International Journal of Animal Bioscience*, 15(7): 100260. DOI: <https://www.doi.org/10.1016/j.animal.2021.100260>
- Farghly MFA, El-Kelawy AMI, Kassab AY, and Emam MS (2021b). Use of palm wastes as alternative flooring materials in broiler chickens houses under prevailing conditions in new valley. *Journal of Desert and Environmental Agriculture*, 1(1): 59-69. DOI: <https://www.doi.org/10.21608/jdea.2021.61866.1006>
- Kralik G, Kralik Z, Kušec ID, Škrtić Z, and Kralik I (2015). Influence of dietary histidine, hybrid line and gender on chicken meat quality and carnosine concentration. *The Journal of Poultry Science*, 52(4): 295-303. DOI: <https://www.doi.org/10.2141/jpsa.0140201>
- Kumar M, Ratwan P, Dahiya SP, and Nehra AK (2021). Climate change and heat stress: Impact on production, reproduction and growth performance of poultry and its mitigation using genetic strategies. *Journal of Thermal Biology*, 97: 102867. DOI: <https://www.doi.org/10.1016/j.jtherbio.2021.102867>
- Lamarca DSF, Pereira DF, Magalhães MM, and Salgado D (2018). Climate Change in Layer Poultry Farming: Impact of heat waves in region of Bastos, Brazil. *Brazilian Journal of Poultry Science*, 20(4): 657-664. DOI: <https://www.doi.org/10.1590/1806-9061-2018-0750>
- Lien R J, Conner DE, and Bilgili SF (1992). The use of recycled paper chips as litter material for rearing broiler chickens. *Poultry Science*, 71(1): 81-87. DOI: <https://www.doi.org/10.3382/ps.0710081>
- Liverpool-Tasie LSO, Sanou A, and Tambo JA (2019). Climate change adaptation among poultry farmers: Evidence from Nigeria. *Climatic Change*, 157: 527-544. DOI: <https://www.doi.org/10.1007/s10584-019-02574-8>
- Manning L, Chadd SA, and Baines RN (2007). Water consumption in broiler chicken: A welfare indicator. *World's Poultry Science Journal*, 63(1): 63-71. DOI: <https://www.doi.org/10.1017/S0043933907001274>
- Mead R (2002). *Statistical methods in agriculture and experimental biology*, 3<sup>rd</sup> Edition. Chapman and

- Hall/CRC., New York, pp. 1-488. DOI: <https://www.doi.org/10.1201/9780203738559>
- Miles DM, Branton SL, and Lott BD (2004). Atmospheric ammonia is detrimental to the performance of modern commercial broilers. *Poultry Science*, 83(10): 1650-1654. DOI: <https://www.doi.org/10.1093/ps/83.10.1650>
- Mohammed HH, Abdelaty AI, Saleem AY, Youssef MI, and Abdel-Hamid SE (2019). Effect of bedding materials on duck's welfare and growth performance. *Slovenian Veterinary Research*, 56(Suppl 22): 149-56. DOI: <https://www.doi.org/10.26873/SVR-752-2019>
- National research council (NRC) (1994). *Nutrient Requirements of Poultry*, 9<sup>th</sup> Revised Edition. The National Academies Press., Washington, DC. Available at: [https://www.agropustaka.id/wp-content/uploads/2020/04/agropustaka.id\\_buku\\_Nutrient-Requirements-of-Poultry\\_Ninth-Revised-Edition-1994-NRC.pdf](https://www.agropustaka.id/wp-content/uploads/2020/04/agropustaka.id_buku_Nutrient-Requirements-of-Poultry_Ninth-Revised-Edition-1994-NRC.pdf)
- Nwaigwe CU, Ihedioha JI, Shoyinka SV, and Nwaigwe CO (2020). Evaluation of the hematological and clinical biochemical markers of stress in broiler chickens. *Veterinary World*, 13(10): 2294-2300. DOI: <https://www.doi.org/10.14202/vetworld.2020.2294-2300>
- Oh SM, Yoon SY, Lee JY, Jeon SM, Oh DY, Ha JJ, Song YH, and Kim JS (2019). Effects of mixed or split-sex feeding on growth performance and behavior of Korean Native Chicken (KNC). *Annals of Animal Resource Sciences*, 30(3): 105-110. DOI: <https://www.doi.org/10.12718/AARS.2019.30.3.105>
- Passini R, De Araujo MAG, De Almeida EA, and Yasuda VM (2012). Evaluation of reflective painting of the roof and artificial ventilation on performance and carcass yield of broilers. *Revista Brasileira de Zootecnia*, 41: 1769-1774. DOI: <http://www.doi.org/10.1590/S1516-35982012000700029>
- Saeed M, Abbas G, Alagawany M, Kamboh AA, Abd El-Hack ME, Khafaga AF, and Chao S (2019). Heat stress management in poultry farms: A comprehensive overview. *Journal of Thermal Biology*, 84: 414-425. DOI: <https://www.doi.org/10.1016/j.jtherbio.2019.07.025>
- Shepherd EM and Fairchild BD (2010). Footpad dermatitis in poultry. *Poultry Science*, 89(10): 2043-2051. DOI: <https://www.doi.org/10.3382/ps.2010-00770>
- Sogunle OM, Egbeyale LT, Bajomo TT, Bamigboje OV, and Fanim AO (2008). Comparison of the performance, carcass characteristics and haematological parameters of broiler chicks reared in cage and floor. *Pakistan Journal of Biological Sciences*, 11(3): 480-483. DOI: <https://www.doi.org/10.3923/pjbs.2008.480.483>
- Thompson HE, Berrang-Ford L, and Ford JD (2010). Climate change and food security in Sub-Saharan Africa: A systematic literature review. *Sustainability*, 2(8): 2719-2733. DOI: <http://www.doi.org/10.3390/su2082719>
- Toghyani M, Gheisari A, Modaresi M, Tabeidian SA, and Toghiani M (2010). Effect of different litter material on performance and behavior of broiler chickens. *Applied animal behaviour science*, 122(1): 48-52. DOI: <https://www.doi.org/10.1016/j.applanim.2009.11.008>
- Topal E and Petek M (2021). Effects of fully or partially slatted flooring designs on the performance, welfare and carcass characteristics of broiler chickens. *British Poultry Science*, 62(6): 804-809. DOI: <https://www.doi.org/10.1080/00071668.2021.1934392>
- Wang Y, Ru YJ, Liu GH, Chang WH, Zhang S, Yan HJ, Zheng AJ, Lou RY, Liu ZY, and Cai HY (2015). Effects of different rearing systems on growth performance, nutrients digestibility, digestive organ weight, carcass traits, and energy utilization in male broiler chickens. *Livestock Science*, 176: 135-140. DOI: <https://www.doi.org/10.1016/j.livsci.2015.03.010>
- Wei FX, Hu XF, Xu B, Zhang MH, Li SY, Sun QY, and Lin P (2015). Ammonia concentration and relative humidity in poultry houses affect the immune response of broilers. *Genetics and Molecular Research*, 14(2): 3160-3169. DOI: <https://www.doi.org/10.4238/2015.April.10.27>
- World organization for animal health (WOAH) (2021). *Terrestrial animal health code. Avian influenza (including infection with high pathogenicity avian influenza viruses)*. Volume I, Paris, France, Chapter 3.3.4, pp. 1-28. Available at: [https://www.woah.org/fileadmin/Home/fr/Health\\_standards/tahm/3.03.04\\_AI.pdf](https://www.woah.org/fileadmin/Home/fr/Health_standards/tahm/3.03.04_AI.pdf)
- World organization for animal health (WOAH) (2021). *Terrestrial animal health code. Newcastle disease (infection with Newcastle disease virus)*. Volume I, Paris, France, Chapter 3.3.14, pp. 1-23. Available at: [https://www.woah.org/fileadmin/Home/fr/Health\\_standards/tahm/3.03.14\\_NEWCASTLE\\_DIS.pdf](https://www.woah.org/fileadmin/Home/fr/Health_standards/tahm/3.03.14_NEWCASTLE_DIS.pdf)
- Yang KY, Ha JJ, Roh H, Cho C, Oh SM, and Oh D (2019). Effects of litter type and gender on behavior characteristics and growth performance of Korean Hanhyup broiler. *Korean Journal of Poultry Science*, 46(3): 155-160. DOI: <http://www.doi.org/10.5536/KJPS.2019.46.3.155>

**Publisher's note:** [Scienceline Publication](#) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access:** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024