

The Interplay of Litter Types, Blends of L-selenomethionine, and Vitamin E in Broiler Chicken's Performance and Health

Eman Mohamed Rady¹ , Gaafar EL-Gendi¹ , Osama El-Garhy¹ , and Hamada Okasha^{1,2*} 

¹Animal Production Department, Faculty of Agriculture, Benha University, Moshtohor 13736, Egypt

²Key Laboratory of Efficient Utilization of Non-grain Feed Resources, College of Animal Science and Technology, Shandong Agricultural University, Taian, Shandong 271018, China

*Corresponding author's E-mail: hamada.okasha@fagr.bu.edu.eg

Received: March 25, 2025, Revised: April 21, 2025, Accepted: May 23, 2025, Published: June 25, 2025

ABSTRACT

The choice of litter type (LT) in broiler chicken production can significantly influence overall performance and health. Adding specific combinations of L-selenomethionine and Vitamin E (L-SeMet + V.E) to drinking water can impact broiler chickens' health. This study aimed to explore the effects of two LT types, wood shavings (WS) and perforated plastic litter (PL) as well as water supplementation with L-SeMet + V.E, on growth performance, blood parameters, antioxidant capacity, and thyroid hormone levels in broiler chickens. A total of 312 one-day-old male broiler chickens were randomly assigned to two different LT groups (WS and PL), with 156 chicks in each. Within each LT group, the chickens were further divided into four subgroups: a control group (T0) that received plain water, and three treatment groups that received drinking water supplements. Each subgroup consisted of three replicates, with 13 broiler chickens per replicate. The water treatments involved varying levels of supplementation with L-SeMet and vitamin E: T1 (0.5 mg/L + 250 mg/L), T2 (1 mg/L + 250 mg/L), and T3 (1.5 mg/L + 250 mg/L). Chickens were assessed for growth performance, feed efficiency, and blood biochemical parameters including lipids, liver enzymes, antioxidants, and thyroid hormones. Chickens raised on PL and receiving T3 exhibited significantly enhanced performance and feed efficiency, outperforming those raised on WS and given different supplementation levels. Significant enhancements were observed in the blood lipid profile and liver enzyme levels across the different LT groups, with the highest values recorded in the PL group. Additionally, broiler chicken in the T2 group, along with the WS × T2 and PL × T3 interactions, showed a notable improvement in blood biochemical parameters. Similarly, chickens raised on PL and given T3-supplemented water, along with the interactions WS × T2 and PL × T3, showed a significant improvement in antioxidant status and thyroid hormone levels in comparison to chickens raised on WS and the other treatment combinations. The use of PL and supplementation with L-SeMet + V.E in drinking water were found to enhance growth performance, improve blood parameters, increase antioxidant capacity, and influence thyroid hormone levels in broiler chickens.

Keywords: Antioxidant, Broiler chicken, L-selenomethionine and Vitamin E, Productivity, Thyroid hormone

INTRODUCTION

To enhance poultry health, it is important to concentrate on both management environment and dietary approaches in order to achieve optimal results. The success of broiler chicken production is influenced by a range of environmental factors, one of the most crucial being the type and proper management of litter material (Pepper and

Dunlop, 2021). The use of high-quality bedding materials in commercial poultry production has attracted considerable attention due to its impact on productive performance (Bilgili et al., 2009). The performance and well-being of broiler chickens is greatly influenced by the type of litter employed in their production (Okasha et al., 2021a; b). It is common knowledge that litter is utilized to

reduce birds' exposure to excrement and to absorb excess moisture (Farghly *et al.*, 2018). However, poor litter quality can lead to the development of pododermatitis in broiler chickens, resulting in discomfort and welfare concerns (Shepherd and Fairchild, 2010), as well as influencing immunological responses, performance, and vulnerability to diseases (De Jong *et al.*, 2014; Wei *et al.*, 2015).

Although litter is traditionally employed in poultry farming, particularly in broiler production, improvements in management practices can enhance both the environmental factors and the well-being of the poultry. One option is to utilize raised plastic flooring, commonly found in broiler chicken barns (Li *et al.*, 2017). Previous study has indicated that broiler chicken productivity can be maintained by utilizing perforated plastic flooring (Almeida *et al.*, 2017), while also improving broiler chicken health (Okasha *et al.*, 2021b). Plastic litter offers durability and cost-effectiveness, as it does not degrade, require replacement, or involve high maintenance costs. It is easy to install and clean plastic litter types, with the potential to reduce occurrences of foot pad injuries (Farghly *et al.*, 2018). Maintaining ideal litter conditions is essential for broiler chicken performance, as excessive moisture, ammonia levels, and the presence of pathogenic organisms can have negative consequences (De Toledo *et al.*, 2020). These include footpad dermatitis, respiratory disorders, and reduced performance such as an unfavorable feed conversion ratio (FCR), decreased body weight (BW), and body weight gain (BWG) (Abougabal and Taboosha, 2023). Therefore, it is essential to meticulously choose and oversee the appropriate litter type in order to ensure optimal broiler chicken health, welfare, and overall performance.

Selenium (Se) is a vital element found in at least 25 selenoproteins that regulate various biological functions in the body (Wang *et al.*, 2017). Selenium's main biological roles include antioxidant defense, adjustment of the immune system, and control of the body's inflammatory reaction (Rayman and Stranges, 2013; Selim *et al.*, 2015). These roles may positively impact the efficiency and metabolic function of broiler chickens. Poultry feed supplements can include both types of selenium, specifically sodium selenate (Na_2SeO_4) and sodium selenite (Na_2SeO_3) (Wickramasuriya *et al.*, 2023). A major advantage of supplying selenomethionine (SeMet) over utilizing inorganic selenium or alternative organic compounds is that SeMet is metabolized with methionine, creating a reserve of selenium in broiler chicken tissues (Surai *et al.*, 2018). According to study of Gangadoo *et al.*

(2020), organic selenium is absorbed and retained more effectively as compared to synthetic forms, with higher concentrations in the duodenum, ileum, and spleen, indicating increased uptake, while lower levels are found in the brain, liver, and breast tissues. To prevent Se deficiency and meet the poultry demands, dietary enrichment with natural selenium is widely practiced due to its superior bioavailability (Surai *et al.*, 2018).

Supplementing broiler chicken diets with selenium has demonstrated favorable impacts on growth, feed markers, immunity, and decreased mortality (Habibian *et al.*, 2016; Calik *et al.*, 2022a). Organic selenium compounds such as hydroxyselenomethionine (OH-SeMet) and L-selenomethionine (L-SeMet) improve selenium reserves in birds, enhance antioxidant protection, and boost resilience to stress in breeder nutrition (Surai and Fisinin, 2014; Wang *et al.*, 2021a; Babazadeh and Ahmadi Simab, 2022). The enhancement of antioxidant capacity has been found to be associated with the activation of selenium-dependent antioxidant enzymes (Liu *et al.*, 2021). In addition, selenium plays a vital role as a component of glutathione peroxidases (Huang *et al.*, 2019). These enzymes, together with vitamin E, contribute to the protection of cells from free radicals (Yoon *et al.*, 2007). Vitamin E is the primary antioxidant that is soluble in lipids, and it holds significant importance in cell membranes (Packer *et al.*, 2001; Gao *et al.*, 2010; Traber and Stevens, 2011). It acts as a crucial lipid antioxidant by interrupting chain reactions and scavenging free radicals within cell membranes and subcellular structures (Young *et al.*, 2003). The studies has shown that adding vitamin E to broiler chicken diets can provide notable benefits due to its potent antimicrobial properties and its capacity to bolster the immune functions (Dalia *et al.*, 2018; Calik *et al.*, 2022a). It can be hypothesized that providing preeminent levels of vitamin E as a supplement can offer benefits in preventing the negative impact of husbandry-related stressors such as transportation, handling, and vaccination, all leading to reduced broiler chicken performance (Perez-Carbajal *et al.*, 2010; Calik *et al.*, 2022b).

The study aimed to evaluate the effects of adding L-selenomethionine and vitamin E (L-SeMet + V.E) supplementation in drinking water on growth performance, feed efficiency, blood profiles, antioxidant enzymes, and T3 and T4 levels. It also explored the impact of this supplementation on broiler chickens raised on either wood shavings or perforated plastic litter, focusing on how it may improve performance and health and mitigate stress effects.

MATERIALS AND METHODS

Ethical approval

The animal care and experimental protocols were approved by the Research Ethics Committee at the Faculty of Agriculture, Moshtohor, Benha University (REC-FOABU/No.16000/3), were observed. The experimental broiler chickens were subjected to the utmost care, and all required actions were implemented to minimize discomfort.

Key features of the different litter types

The study utilized a washable perforated plastic floor (Al-Ekhlash for Plastics Industry and Trading, Qalyubia, Egypt), measuring $100 \times 100 \times 15$ mm (length \times width \times height), and weighing 1.75 kg with a resistance of up to 200 kg/m² per unit. The wood shavings (beech) used for the poultry litter were obtained from a nearby tree processing facility. These soft, dry, and non-toxic shavings were spread at a depth of 7 cm. The materials exhibited high absorbency and were resistant to clumping.

Animals and study design

All broiler chicks (Ross 308) were obtained from a commercial hatchery (Association of Al-Tanmia for Hatching and Poultry Production, Qalyubia, Egypt). The chicks were transported in a temperature-controlled vehicle for about one hour to the experimental farm located at the Poultry Research Unit, Faculty of Agriculture, Benha University, Egypt. Upon arrival, the chicks were housed in research enclosures within a controlled-geographical area, where stringent measures were implemented to maintain optimal experimental and hygienic conditions. All pens, each measuring 1×1 meter, were placed in the same location to ensure consistency. The stocking density in each pen was maintained at 13 chicks per square meter. Continuous electric lighting was provided for the first five days, after which the lighting schedule was adjusted to 20 hours of light and 4 hours of darkness for the following 30 days. Lighting intensity was applied in accordance with Directive 2007/43/EC in order to enhance the welfare of broiler chickens. Gas-powered space heaters were used in this experiment. The air temperature and relative humidity were kept at 33°C and 63%, respectively, for as long as three days before the chicks were placed in the pens. Throughout the initial week of the trial, the temperature was maintained at 33°C. Following this, the temperature was gradually reduced by 2-3°C per week until it reached the target range of 25°C by day 28. The broiler chickens were vaccinated according to a specified schedule: On day 8, they were administered the

Newcastle Disease Virus (NDV) Clone 30 vaccine via eye drop, in addition to a bivalent NDV vaccine administered through injection under the neck membrane. The Gumboro intermediate vaccine was given at 12 days of age, followed by the NDV Clone 30 vaccine again at 21 days of age. The inactivated Avian Influenza (AI) H5N2 vaccine was administered at 10 days of age (Nobilis, Intervet, Boxmeer, The Netherlands). The immunization process followed established guidelines and protocols, and a licensed veterinarian supervised its implementation.

For a 35-day experiment, a total of 312 one-day-old male Ross 308 broiler chicks with an average initial body weight of 46.22 ± 0.1 g were used. The study employed a 2×4 factorial arrangement of treatments with three replicates, resulting in 24 experimental groups of chickens ($n = 13$ per group). Each group served as an experimental unit for the study. The chicks were randomly divided into two primary groups based on litter types: wood shavings and perforated plastic litter ($n = 156$ chicks per litter type). The chicks from each litter type were randomly allocated into a control group and three experimental groups ($n = 39$ chicks per group), which received different levels of L-SeMet + V.E into their drinking water. The experimental sets were labeled as follows: T1, (0.5 mg L-SeMet + 250 mg V.E); T2, (1 mg L-SeMet + 250 mg V.E); and T3, (1.5 mg L-SeMet + 250 mg V.E) mg/L, respectively. In this study, the supplementation level of vitamin E was increased in alignment with the guidelines provided by Aviagen (Aviagen, 2019). This finding aligns with earlier study, where an average supplementation of 200 mg/kg of vitamin E was shown to enhance oxidative status, improve performance parameters, and positively influence meat quality in broiler chickens (Mazur-Kušnerek et al., 2019). In the current study, the organic selenium (Se) content in the diet was adjusted to the maximum permissible level, ranging from 1 to 3 mg/kg, consistent with the levels utilized in prior studies (Calik et al., 2022b; Kazem and Al-Khafaji, 2023). The vitamin E product added to the water was a misallied liquid solution that could disperse in water. It contained 250 mg of α -tocopherol per liter (L-selenomethionine® and vitamin E®, Alcon Biosciences Pvt. Ltd, Mumbai, India). Table 1 details the nutritional makeup of the experimental diets meticulously computed and analyzed, and then utilized in various stages of the experiment. The broiler chickens were fed a pelleted basal diet formulated to meet and slightly exceed the nutrient requirements specified by the National Research Council (NRC, 1994). Both feed and water were provided *ad libitum* for the entire duration of the study.

Table 1. The composition of ingredients and nutrients in the basal starter and grower diets in broiler chickens (Ross 308)

Ingredients (%)	Starter diet (0-3 weeks)	Grower diet (3-5 weeks)
Yellow Corn	60.56	65.4
Soybean meal (40%)	22	16.1
Corn gluten meal (60%)	12.6	13.2
Calcium hydrogen phosphate	1.5	1.7
Limestone	2.01	2.32
Salt (NaCl)	0.42	0.42
VIT and Min premix ¹	0.3	0.3
Choline chloride (50%)	0.2	0.2
L-lysine (78%)	0.24	0.24
Methionine	0.17	0.12
Total	100	100
Energy and nutrient composition		
ME (kcal/kg)	3000	3053
Crude protein (%)	22.29	20.77
EE (%)	2.80	3.00
Crude fiber, %	3.00	2.77
Ca (%)	1.21	1.19
T. phosphorus (%)	0.65	0.62
Av. phosphorus (%)	0.42	0.40
Lys (%)	1.11	0.95
Met (%)	0.50	0.40
Met + Cys (%)	0.95	0.80

¹This feed formulation comprises a mineral and vitamin premix, providing the following concentrations (mg/3kg): Iron 80, Manganese 100, Copper 8, Zinc 75, Selenium 0.15, and Iodine 0.35. Additionally, it includes vitamin A (12,500 IU), vitamin D3 (2,500 IU), vitamin E (30 IU), vitamin K3 (2.65 mg), vitamin B1 (2 mg), vitamin B2 (6 mg), vitamin B3 (50 mg), vitamin B5 (12 mg), vitamin B7 (0.0325 mg), vitamin B9 (1.25 mg), and vitamin B12 (0.025 mg) per kilogram. These values were calculated based on feedstuff data as shown in the NRC (1994). Abbreviations: ME (kcal/kg): Metabolisable Energy (kcal/kg), EE (%): Ether Extract (%), T. phosphorus (%): Total phosphorus (%), Av. phosphorus (%): Available phosphorus (%), Met (%): Methionine, Met + Cys (%) Methionine and Cystine (%)

Taking samples and measurements

Growth performance and feed efficiency

Broiler chickens were weighed weekly to individually assess their growth performance, including body weight (BW) and body weight gain (BWG). Additionally, daily feed intake (FI) was recorded in replicates to calculate the ratio between (FI/g: BWG/g) expressed as feed conversion efficiency ratio (FCR). The European Production Efficiency Factor (EPEF) was calculated using the following equation: $EPEF = (Livability[\%] \times BW [kg]) / (FCR \times Trial\ duration) \times 100$.

Blood samples

Twelve broiler chickens were chosen at random from each main group at the end of the 35-day trial (four per replicate). After a 12-hour period of fasting, 5 mL of blood was extracted from the wing vein. The samples were

carefully placed into sterile tubes containing heparin and serum separator for subsequent analysis of thyroid hormones. The tubes were centrifuged for ten minutes at 4°C at 3500 × g to isolate the plasma. The plasma lipid profile, including total lipids, cholesterol (CHO), triglycerides (TRIG), low-density lipoproteins (LDL), and high-density lipoproteins (HDL), was analyzed using commercially available kits (Meikang Incorporated, Ningbo, China) following the methodology outlined by Gornall et al. (1949). Additionally, liver enzymes such as aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were measured using a colorimetric method as described by Sirois (2019). The enzymes were measured using kits procured from the Nanjing Jiancheng Institute of Bioengineering (Jiangsu, China).

Evaluating antioxidant capacity

To assess the concentration of total antioxidants (T-AOC) in plasma, a commercially available kit (Randox, UK) was utilized, which was based on the method developed by Miller et al. (1993). The activity of blood glutathione peroxidase (GPx, EC 1.11.1.9) was assessed using commercially available GPx kits from Randox (Crumlin, UK), adhering to the manufacturer's instructions. The method for GPx measurement was based on the procedure established by Paglia and Valentine (1967). Additionally, superoxide dismutase (SOD, EC 1.1.5.1) activity in erythrocyte lysates was measured using kits provided by Randox Laboratories Ltd. (Crumlin, UK), following the protocols outlined by Woolliams et al. (1983). Moreover, plasma malondialdehyde (MDA) levels were measured using a modified fluorometric technique as detailed by Jo and Ahn (1998).

Measuring thyroid hormone levels

Serum samples were used to assess the levels of thyroid hormones (T3 and T4) using radioimmunoassay (RIA) kits, following the manufacturer's instructions. The techniques outlined by Renden et al. (1994) were employed for this analysis.

Statistical analysis

The pen served as the experimental unit (n = 24), and group means were subjected to statistical analysis. The data underwent analysis utilizing the General Linear Models (GLM) procedure of two-way ANOVA in SPSS (IBM SPSS, Version 27.0. Armonk, NY). Tukey's test was utilized to differentiate between treatments and assess the differences. The linear model was specified as:

$$Y_{ijk} = \mu + LT_i + T_j + (LTT)_{ij} + e_{ijk}$$

where Y_{ijk} represents the k th observation, and μ stands for the overall mean. The term LT_i refers to the effect of the i th litter type (wood shavings and plastic litter), whereas T_j represents the effect of the j th L-SeMet + V.E treatment level (0, 0.5, 1, 1.5 + 250 V.E per mg/L). Additionally, the term $(LTT)_{ij}$ represents the interaction effect between the i th litter type and the j th L-SeMet + V.E treatment. The term e_{ijk} refers to the experimental error, assuming a mean of zero ($\bar{X} = 0$) and a variance of σ^2_e . Statistical significance was assessed using a P-value threshold of 0.001, unless otherwise specified.

RESULTS

Growth performance and feed efficiency

Table 2 highlights the effects of litter type and L-SeMet + V.E supplementation on broiler chickens' growth and feed efficiency up to 35 days. Significant improvements ($p < 0.001$) were observed in growth performance (BW and BWG) and feed efficiency (CFI, CFCR, and EPEF) due to both litter type and L-SeMet + V.E inclusion. While initial body weights showed no differences ($p = 0.648$), supplementation had a notable impact on growth traits during later periods. The addition of 1.5 + 250 mg/l L-SeMet + V.E to the drinking water resulted in a significant increase in BW, BWG EPEF and decreased CFCR ($p < 0.001$). Interactions between WS with T2 and PL with T3 showed the most significant improvements in growth, feed efficiency, and European Production Efficiency Factor (EPEF) ($p < 0.001$).

Blood biochemical analysis

Table 3 summarizes the findings of the blood lipid profile and liver enzyme analysis. Statistically significant differences ($p < 0.001$) were detected in the plasma concentrations of CHO, TRIG, LDL, HDL, and AST among broiler chickens reared on WS and PL litters. The incorporation of 1 + 250 mg/l L-SeMet + V.E into the drinking water significantly altered all blood parameters ($p < 0.001$), with increased HDL levels and reduced lipid profiles and liver enzymes. Some interactions such as WS \times T2 increased HDL, while others such as PL \times T3 reduced CHO, TRIG, and LDL levels. Both PL \times T3 and WS \times T2 significantly decreased AST and ALT levels, respectively, as compared to other interactions ($p < 0.001$).

Antioxidants and thyroid hormones

Table 4 and Graph 1 show significant differences ($p < 0.001$) in plasma antioxidants (TAOC, GPX, MDA) and serum thyroid hormones (T3, T4) among broiler chickens raised on different litter types. Chickens on PL litter exhibited higher antioxidant levels and thyroid hormones with lower MDA, as compared to those on WS litter. Supplementation with L-SeMet + V.E at 1.5 + 250 mg/L significantly improved antioxidant status and thyroid hormone levels across all parameters ($p < 0.001$). Interaction analysis revealed that chickens raised on both litter types and given T2 or T3 had the highest antioxidant capacity (TAOC, GPX, SOD) and thyroid hormones (T3, T4), with the lowest MDA levels ($p < 0.001$).

Table 2. Effects of different litter types, L-Selenomethionine + Vitamin E, and their interactions on broiler chickens' growth and feed efficiency

Items		BW (g)		BWG (g)	CFI (g/bird)	CFCR	EPEF
		0	5 WKS	0-5 WKS	0-5 WKS	0-5 WKS	0-5 WKS
Litter types (LT)	WS	46.26	2175.47 ^b	2129.05 ^b	3094.50 ^b	1.46 ^a	425.72 ^b
	PL	46.49	2307.66 ^a	2261.48 ^a	3137.67 ^a	1.39 ^b	453.72 ^a
	SEM	0.342	3.930	16.604	6.057	0.014	6.593
L-SeMet + V.E (T) mg/l	T0	47.22	2218.66 ^c	2171.44 ^b	3132.79 ^b	1.44 ^b	428.44 ^b
	T1	46.34	2048.00 ^d	2001.65 ^c	3072.73 ^c	1.53 ^a	398.52 ^c
	T2	46.29	2282.52 ^b	2236.23 ^b	3100.05 ^{bc}	1.38 ^{bc}	464.30 ^a
	T3	45.66	2417.09 ^a	2371.74 ^a	3165.80 ^a	1.35 ^c	484.84 ^a
	SEM	0.484	5.558	23.481	9.464	0.20	8.847
LT \times T	WS \times T0	47.56	2149.33 ^e	2101.77 ^{cde}	3156.73 ^b	1.50 ^{ab}	408.86 ^{cd}
	WS \times T1	45.86	2084.00 ^f	2038.14 ^{de}	3050.87 ^d	1.50 ^{ab}	406.77 ^{cd}
	WS \times T2	46.52	2335.71 ^b	2289.17 ^b	3090.50 ^{cd}	1.35 ^{cd}	483.32 ^b
	WS \times T3	46.00	2132.86 ^e	2087.14 ^{cde}	3104.93 ^{bcd}	1.49 ^{abc}	409.63 ^{cd}
	PL \times T0	46.88	2288.00 ^c	2241.12 ^{bc}	3104.53 ^{bcd}	1.39 ^{bc}	451.28 ^{bc}
	PL \times T1	46.83	2012.00 ^g	1965.17 ^e	3094.60 ^{cd}	1.57 ^a	390.27 ^d
	PL \times T2	46.05	2229.33 ^d	2183.29 ^{bcd}	3109.60 ^{bcd}	1.42 ^{bc}	445.23 ^{bc}
	PL \times T3	45.31	2701.33 ^a	2656.35 ^a	3226.67 ^a	1.21 ^d	560.05 ^a
	SEM	0.684	7.861	33.207	13.382	0.280	10.339
p-value	LT	0.648	<.001	<.001	<.001	0.007	<.001
	T	0.193	<.001	<.001	<.001	<.001	<.001
	LT \times T	0.572	<.001	<.001	<.001	<.001	<.001

The data sets reflect the mean values obtained from three replications per treatment. Different letters denote significant differences ($p < 0.001$) among means within the same column for each factor and interactions. Abbreviations: BW: Body weight, BWG: Body weight gain, CFI: Cumulative feed intake, CFCR: Cumulative feed conversion ratio, EPEF: European production efficiency factor, L: Litter types, WS: Wood shavings, PL: Perforated plastic litter, T: Treatments, WKS: Weeks, L-SeMet: L-selenomethionine, V.E: Vitamin E, SEM: Standard error mean, T0: Control group (plain water), T1: 0.5 + 250 mg/l L-SeMet + V.E, T2: 1 + 250 mg/l L-SeMet + V.E, T3: 1.5 + 250 mg/l L-SeMet + V.E

Table 3. Effects of different litter types, L-Selenomethionine + Vitamin E, and their interactions on broiler chickens' lipid profile and liver enzymes


Items		TL (mg/dl)	CHO (mg/dl)	TRIG (mg/dl)	LDL (mg/dl)	HDL (mg/dl)	AST (U/l)	ALT (U/l)
Litter Types (LT)	WS	851.98	216.66 ^a	74.07 ^a	156.92 ^a	44.92	152.64 ^a	57.96
	PL	779.76	123.21 ^b	58.02 ^b	70.14 ^b	41.46	144.40 ^b	58.29
	SEM	7.208	2.699	2.430	2.083	0.768	1.067	1.578
L-SeMet + V.E (T) mg/l	T0	1130.95 ^a	198.81 ^a	101.23 ^a	140.56 ^a	35.06 ^b	1.66.79 ^a	72.42 ^a
	T1	821.42 ^b	173.81 ^{ab}	77.775 ^b	112.70 ^b	39.47 ^b	143.59 ^b	57.11 ^b
	T2	549.21 ^c	146.43 ^c	42.590 ^c	98.445 ^d	49.75 ^a	141.26 ^b	44.91 ^c
	T3	761.90 ^b	160.71 ^{bc}	42.595 ^c	102.44 ^{bc}	48.49 ^a	142.45 ^b	58.06 ^b
	SEM	10.312	3.862	3.436	2.981	1.084	1.526	2.231
LT × T	WS × T0	1071.43 ^{ab}	242.86 ^a	106.17 ^a	183.42 ^a	32.33 ^d	178.52 ^a	74.30 ^a
	WS × T1	857.14 ^{bcd}	228.57 ^a	87.65 ^{ab}	160.66 ^{ab}	38.21 ^{cd}	140.28 ^{bcd}	44.63 ^b
	WS × T2	550.81 ^e	176.19 ^b	49.38 ^{cd}	133.24 ^b	58.78 ^a	139.26 ^{cd}	42.56 ^b
	WS × T3	928.57 ^{abc}	219.05 ^a	53.09 ^{cd}	150.39 ^b	50.38 ^{ab}	152.52 ^{bc}	70.36 ^a
	PL × T0	1190.48 ^a	154.76 ^{bc}	96.30 ^a	97.71 ^c	37.79 ^{cd}	155.07 ^b	70.54 ^a
	PL × T1	785.71 ^{cde}	119.05 ^{cd}	67.90 ^{bc}	64.74 ^d	40.73 ^{bcd}	146.91 ^{bcd}	69.60 ^a
	PL × T2	547.62 ^e	116.67 ^{cd}	35.80 ^d	63.65 ^d	40.73 ^{bcd}	143.26 ^{bcd}	47.26 ^b
	PL × T3	595.24 ^{de}	102.38 ^d	32.10 ^d	54.94 ^d	46.60 ^{bc}	132.38 ^d	45.76 ^b
	SEM	14.583	5.461	4.860	4.215	1.533	2.16	3.156
Two-way ANOVA								
p-value	LT	0.086	<.001	<.001	<.001	0.054	0.001	0.885
	T	<.001	<.001	<.001	<.001	<.001	<.001	<.001
	LT × T	0.007	0.01	<.001	0.054	<.001	<.001	<.001

The data sets reflect the mean values obtained from three replications (4 chickens in each replicate) per treatment. Different letters denote significant differences ($p < 0.001$) among means within the same column for each factor and interactions. Abbreviations: L: Litter types, WS: Wood shavings, PL: Perforated plastic litter, T: Treatments, L-SeMET: L-selenomethionine, V.E: Vitamin E, T0: Control group (plain water), T1: 0.5 + 250 mg/l L-SeMET + V.E, T2: 1 + 250 mg/l L-SeMET + V.E, T3: 1.5 + 250 mg/l L-SeMET + V.E, SEM: Standard error mean, TL: Total lipids, CHO: Cholesterol, TRIG: Triglycerides, LDL: Low density lipoproteins, HDL: High density lipoproteins, AST: Aspartate aminotransferase, ALT: Alanine aminotransferase

Table 4. Effects of different litter types, L-Selenomethionine + Vitamin E, and their interactions on broiler chickens' antioxidant levels

Items		TAOC (mM/L)	GPX (U/GT)	SOD (U/ml)	MDA (nmol/mL)
Litter Types (LT)	WS	0.77 ^b	43.67 ^b	3.48	3.65 ^a
	PL	0.89 ^a	48.62 ^a	3.52	2.37 ^b
	SEM	0.011	0.452	0.112	0.052
L-SeMet + V.E (T) mg/l	T0	0.74 ^b	19.21 ^d	2.81 ^b	5.28 ^a
	T1	0.77 ^b	29.22 ^c	3.27 ^b	3.25 ^b
	T2	0.83 ^b	58.35 ^b	3.94 ^a	2.05 ^c
	T3	0.97 ^a	77.81 ^a	3.99 ^a	1.43 ^c
	SEM	0.016	0.414	0.158	0.074
LT × T	WS × T0	0.67 ^c	19.45 ^e	2.88 ^c	6.30 ^a
	WS × T1	0.77 ^{bc}	38.90 ^d	3.30 ^{bc}	1.96 ^c
	WS × T2	0.88 ^b	97.26 ^a	4.38 ^{ab}	1.15 ^c
	WS × T3	0.77 ^{bc}	38.90 ^d	3.38 ^{bc}	5.20 ^{ab}
	PL × T0	0.82 ^{bc}	18.98 ^e	2.75 ^c	4.26 ^b
	PL × T1	0.77 ^{bc}	19.55 ^e	3.25 ^c	2.21 ^c
	PL × T2	0.79 ^{bc}	58.36 ^c	3.50 ^{bc}	1.72 ^c
	PL × T3	1.18 ^a	77.80 ^b	4.60 ^a	1.31 ^c
	SEM	0.023	0.586	0.224	0.105
Two-way ANOVA					
p-value	LT	<.001	<.001	0.804	<.001
	T	<.001	<.001	<.001	<.001
	LT × T	<.001	<.001	0.002	<.001

The data sets reflect the mean values obtained from three replications (4 chickens in each replicate) per treatment. Different letters denote significant differences ($p < 0.001$) among means within the same column for each factor and interactions. Abbreviations: L: Litter types, WS: Wood shavings, PL: Perforated plastic litter, T: Treatments, L-SeMET: L-selenomethionine, V.E: Vitamin E, SEM: standard error mean, T0: Control group (plain water), T1: 0.5 + 250 mg/l L-SeMET + V.E, T2: 1 + 250 mg/l L-SeMET + V.E, T3: 1.5 + 250 mg/l L-SeMet + V.E, TAOC: Total antioxidants (mM/L), GPX: Glutathione peroxidase (U/GT), SOD: Superoxide dismutase (U/ml), MDA: Malondialdehyde (nmol/ml)



Graph 1. Effects of litter types, L-Selenomethionine + Vitamin E, and their interactions on serum levels of T3 (A) and T4 (B) in broiler chickens. The data represent mean values from three replicates (4 chickens in each replicate) per treatment. The use of distinct letters placed above the columns indicates significant variances, established by Tukey's test with a significance level of $p < 0.001$. Abbreviations: L: Litter types, WS: Wood shavings, PL: Perforated plastic litter, T: Treatments, L-SeMET: L-selenomethionine, V.E: Vitamin E, T0: Control group (plain water), T1: 0.5 + 250 mg/l L-SeMET + V.E, T2: 1 + 250 mg/l L-SeMET + V.E, T3: 1.5 + 250 mg/l L-SeMET + V.E, T3: Triiodothyronine hormone (ng/dl), T4: Thyroxin hormone (µg/dl)

DISCUSSION

Growth performance and feed efficiency

There is a notable emphasis on enhancing the productive performance of broiler chickens. Achieving better performance requires the implementation of efficient management practices, ensuring optimal housing conditions, and carefully formulating their diet (Bist et al., 2024). The findings of the current investigation unveiled that broiler chickens raised on PL litter type exhibited significantly superior growth performance compared to those housed on other WS litter systems. This was evident in parameters of growth performance (BW and BWG), and feed efficiency (CFI, FCR, and EPEF). These results are attributed to the fact that broiler chickens reared on perforated plastic litter types consumed a higher amount of feed. This is likely due to a reduced likelihood of pecking in the litter, which leads to increased feed intake. Consequently, this positively affects body weight (BW) and leads to an increase in body weight gain (BWG). Chickens raised on plastic flooring exhibited comparatively greater weight gains but had a lower FCR compared to chickens reared solely on wood shavings (Almeida et al., 2017; Çavuşoğlu et al., 2018). This improvement may be attributed to the fact that WS bedding absorbs broiler chicken feces, resulting in

increased moisture and interaction with ammonia. Consequently, the surface temperature of the bedding is somewhat elevated, which affects the environmental comfort for the broiler chickens. This leads to a decrease in feed intake and adversely affects feed conversion efficiency. However, this is not the case with PL litter type, where broiler chickens kept in deep litter systems exhibited the lowest FI and BW (Abd El-Wahab et al., 2020). Broiler chickens raised on plastic litter exhibit a lower pecking at litter, which results in increased feed intake. The higher observed feed intake could account for the variations in growth performance and feed efficiency among different litter types. Reducing contact with fecal matter improves the overall health of broiler chickens, leading to enhanced growth performance (Wang et al., 2015). In the present study, broiler chickens reared on PL litter exhibited significantly higher values for body weight gain, feed intake, and feed conversion ratio compared to those raised on WS litter. Without a doubt, the factors encompassed within the European Production Efficiency Factor (EPEF), including bird livability percentage, body weight (kg), FCR, and days until slaughter, as discussed and examined earlier, demonstrate superior performance in broiler chickens housed in PL systems, resulting in a higher EPEF score.

The findings of this study show that supplementing broiler chickens' drinking water with organic L-SeMet combined with vitamin E is more effective in preserving growth performance and improving feed efficiency (Table 2). The higher supplementation rates (1 and 1.5 mg/L L-SeMet + 250 mg/L V.E) yielded more pronounced positive effects. This finding indicates that the combination of organic L-SeMet was effective in enhancing energy utilization of feed. The findings from this study align with the results of [Calik et al. \(2022b\)](#), suggesting that incorporating organic selenium + V.E into broiler chickens' diet at a level of 1:250 mg/kg can significantly enhance both BWG and FI. The observed enhancement in productive performance in this investigation can be attributed to antioxidant properties of organic L-SeMet and V.E, which help protect against potential stressors encountered during standard husbandry practices. Selenium plays crucial roles in the body, primarily as a key component of selenoproteins ([Wang et al., 2021a](#)). These selenoproteins participate in various functions, comprising the metabolism of thyroid hormones. Specifically, they facilitate the transformation of T4 into its active state T3 ([Schomburg, 2012](#)), which has a crucial function in controlling energy and protein assimilation in the body ([Wickramasuriya et al., 2023](#)). The current study suggests that increased thyroid hormone levels significantly enhance growth performance and feed efficiency in broiler chickens. However, limited published studies exist on the effects of L-SeMet + V.E supplementation via drinking water under standard husbandry practices, indicating a need for further investigation.

Blood biochemical analysis

The assessment of blood biochemical factors serves as a valuable tool for identifying metabolic and nutritional changes, which are crucial for assessing the overall health condition in poultry ([Ghasemi et al., 2013](#)). The study revealed significant effects of litter type, L-SeMet + V.E supplementation, and their interaction on blood biochemical factors. Broiler chickens raised on both litter types and given T2 or T3 showed increased HDL levels and reduced plasma lipids and liver enzymes. The findings of the current study are in line with a former experiment conducted by [Okasha et al. \(2021b\)](#) on broiler chickens. Similarly, broiler chickens reared on wood shavings and sand showed significantly lower levels of CHO, TRIG, LDL compared to those reared on plastic litter. Additionally, higher levels of HDL were observed in broiler chickens reared on WS litter compared to those

raised on sand litter ([Lasheen et al., 2023](#)). The study found higher AST activity in broiler chickens raised on wood shavings compared to perforated plastic litter, though it did not impact broiler chickens' well-being. These results are consistent with previous findings, with AST activity highest in the avian heart, followed by the liver and skeletal muscle ([Ognik and Krauze, 2016](#); [Costa et al., 2021](#)).

Incorporation of L-SeMet + V.E into drinking water led to decreased plasma lipid profile and liver enzymes, alongside increased plasma HDL levels in broiler chickens raised on both types of litter, as indicated in Table 3. These findings align with [Kang et al. \(2000\)](#) who proposed that selenium has a cholesterol-lowering effect, decreasing both cholesterol and triglycerides levels. Additionally, [Amer et al. \(2019\)](#) discovered that incorporating Se, irrespective of its origin, resulted in a reduction in TRIG levels in rabbits. Selenium plays a role in metabolism of lipids ([Zhang et al., 2018](#)). The study suggests that L-SeMet + V.E supplementation improves lipid metabolism by increasing HDL levels and reducing LDL levels, which is beneficial as lower LDL minimizes the risk of oxidase-low-density lipoproteins (ox-LDL) formation and cholesterol buildup in arterial walls ([Toth, 2005](#)). Therefore, selenium has a crucial function in safeguarding cells against oxidative stress by boosting the activity of glutathione peroxidase ([Negis et al., 2006](#)). Incorporation of L-SeMet + V.E had a significant impact on the serum ALT and AST levels in broiler chickens. This is reinforced by the discovery of [Dalia et al. \(2017\)](#), who observed a notable reduction in the liver enzymes in chickens fed organic selenium. Moreover, [Biswas et al. \(2010\)](#) documented comparable results, noting a reduction in ALT and AST concentrations in broiler chickens that received a diet enriched with Se at a level of 0.5 and 1 mg/kg. A decrease in lipid profile and liver enzyme levels indicates improved protection against oxidative damage by enhancing the redox state. These findings are supported by the antioxidant status results obtained in this study.

Antioxidants and thyroid hormones

In avian physiology, monitoring oxidative status plays a crucial role in assessing health changes ([Surai et al., 2019](#)). Assessing antioxidant levels in broiler chickens typically involves analyzing parameters such as T-AOC and MDA in the bloodstream, as well as the activity of enzymatic scavengers SOD and GPx ([Ghasemi et al., 2020](#)). Broiler chickens raised on PL litter and supplemented with 1.5 mg/L of L-SeMet + V.E showed improved serum antioxidant status and thyroid hormones,

highlighting its protective role against oxidative stress. In line with the present results, [Ghanima et al. \(2020\)](#) and [El-Maaty et al. \(2023\)](#) demonstrated that broiler chickens reared on perforated PL exhibited improved antioxidant status in contrast to those reared on floor litter or in batteries. According to [Okasha et al. \(2021b\)](#), substantial differences were noted in the plasma GPX and MDA levels of broiler chickens as a result of the litter type. Antioxidant and thyroid hormone levels in broiler chickens raised on wood shavings may be influenced by stress factors such as mold, iodine interference, and ammonia exposure ([Şen et al., 2023](#)). Further research is needed to fully understand the effects of different litter types on broiler chicken production under standard practices.

Chickens' antioxidant system is regulated by numerous crucial enzymes, comprising T-AOC, GPx, SOD, and MDA ([Wang et al., 2021b](#)). Antioxidant properties in selenium stem from its role in GSH-Px, where it forms part of the enzyme's active center, aiding in the removal of unstable molecules generated during metabolism ([Huang et al., 2019](#); [Deng et al., 2022](#)). The findings suggest reduced lipid oxidation, likely due to the higher bioavailability of organic selenium combined with V.E., which enhances selenium retention in the body. Consequently, this increased retention contributes to elevated levels of GSH-Px ([Zhang et al., 2014](#)). Likewise, [Dalia et al. \(2017\)](#) found that bacterial organic selenium (BOSe) is a superior selenium source for broiler chickens, enhancing antioxidant levels and glutathione peroxidase gene activation more effectively than inorganic selenium. Thyroid hormones (T3 and T4) play a vital role in gluconeogenesis and glycogen synthesis in poultry, facilitating the transformation of non-carbohydrate substrates into glucose and promoting the production and storage of glycogen in the liver ([Duntas and Brenta, 2018](#)). The current study revealed that L-SeMet + V.E supplementation elevated T3 and T4 levels in broiler chickens, likely attributed to the functions of selenium in promoting the conversion of T4 to T3 and sustaining selenium proteins. Similarly, [Lin et al. \(2014\)](#) demonstrated that selenium deficiency hinders the conversion of T4 to T3 in chickens by decreasing the activity of critical metabolic enzymes (Iodothyronine Deiodinase Type-1, Iodothyronine Deiodinase Type-2, and Iodothyronine Deiodinase Type-3) and selenoproteins (Thioredoxin Reductase 2, Selenoprotein I, Selenoprotein U, Glutathione peroxidase 1, and Glutathione Peroxidases 2), thereby indirectly disrupting thyroid hormone metabolism. Enhancing antioxidant response and

regulating T3 and T4 through nutrition is of growing interest. The study hypothesizes that supplementing L-SeMet and Vitamin E in drinking water may improve antioxidant and thyroid hormone levels, boosting broiler chickens' performance and feed efficiency. Supplementing L-SeMet and Vitamin E in drinking water could lead to cost savings through improved feed efficiency, reduced health issues, and enhanced broiler chickens' welfare due to improved health and growth. These benefits make it a promising strategy for poultry farming.

CONCLUSION

Broiler chickens raised on perforated plastic litter and supplemented with L-SeMet + Vitamin E in drinking water showed improved growth, higher feed efficiency, enhanced antioxidant status, and elevated T3 and T4 levels. The supplementation also improved blood parameters, increasing HDL and reducing lipid profile and liver enzyme levels. This approach demonstrates significant benefits for broiler chicken productivity and health under standard husbandry practices. By enhancing strategies for organic supplementation, assessing their long-term impacts, and investigating the interactions between L-SeMet and other nutrients, broiler chicken farming practices could be significantly advanced. These initiatives would contribute to sustainable, efficient, and welfare-focused poultry production in the future.

DECLARATIONS

Acknowledgments

The authors express their appreciation to the Faculty of Agriculture at Benha University for their authorized ethical assistance in conducting this study.

Authors' contributions

Hamada Okasha, Eman Rady, Osama El-Garhy, and Gaafar EL-Gendi authored the original text, contributed to the experimental design, carried out the experiments, and conducted the statistical analysis. Hamada Okasha and Eman Rady contributed to both the statistical analysis and the design of the experiments. Hamada Okasha, Eman Rady, Osama El-Garhy, and Gaafar EL-Gendi analyzed and reviewed the results and then composed the final essay. All authors have reviewed and approved the final version of the manuscript for publication.

Competing interests

The authors declare no conflicts of interest.

Ethical considerations

Ethical issues, including plagiarism, consent to publish, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy, have been checked by all the authors.

Availability of data and materials

The data obtained in this study can be obtained from the corresponding author upon reasonable request.

Funding

This study was not supported by any external funding sources.

REFERENCES

- Abd El-Wahab A, Kriewitz JP, Hankel J, Chuppava B, Ratert C, Taube V, Visscher C, and Kamphues J (2020). The effects of feed particle size and floor type on the growth performance, GIT development, and pododermatitis in broiler chickens. *Animals*, 10(8): 1256. DOI: <http://www.doi.org/10.3390/ani10081256>
- Abougabal MS and Taboosha M F (2023). Effect of different types of available litter materials on the performance and welfare of broiler chickens. *Al-Azhar Journal of Agricultural Research*, 48(1): 139-152. DOI: <https://www.doi.org/10.21608/ajar.2023.316064>
- Almeida DE, Eduardo Alves, de Souza LFA, Sant'Anna AC, Bahiense RN, Macari M, and Furlan RL (2017). Poultry rearing on perforated plastic floors and the effect on air quality, growth performance, and carcass injuries—Experiment I: Thermal comfort. *Poultry Science*, 96(9): 3155-3162. DOI: <https://www.doi.org/10.3382/ps/pex131>
- Amer SA, Omar AE, and Abd El-Hack ME (2019). Effects of selenium- and chromium-enriched diets on growth performance, lipid profile, and mineral concentration in different tissues of growing rabbits. *Biological Trace Element Research*, 187: 92-99. DOI: <https://www.doi.org/10.1007/s12011-018-1356-4>
- Aviagen (2019). Ross 308 broiler: Nutrition specifications. Available at: https://eu.aviagen.com/assets/Tech_Center/Ross_Broiler/RossBroilerNutritionSpecs2019-EN.pdf
- Babazadeh D and Ahmadi Simab P (2022). Methionine in poultry nutrition: A review. *Journal of World's Poultry Science*, 1(1): 1-11. DOI: <https://www.doi.org/10.58803/jwps.v1i1.1>
- Bilgili S, Hess J, Blake J, Macklin K, Saenmahayak B, and Sibley J (2009). Influence of bedding material on footpad dermatitis in broiler chickens. *Journal of Applied Poultry Research*, 18(3): 583-589. DOI: <https://www.doi.org/10.3382/japr.2009-00023>
- Bist RB, Bist K, Poudel S, Subedi D, Yang X, Paneru B, Mani S, Wang D, and Chai L (2024). Sustainable poultry farming practices: A critical review of current strategies and future prospects. *Poultry Science*, 103(12): 104295. DOI: <https://www.doi.org/10.1016/j.psj.2024.104295>
- Biswas A, Ahmed M, Bharti V, and Singh S (2010). Effect of antioxidants on physio-biochemical and hematological parameters in broiler chicken at high altitude. *Asian-Australasian Journal of Animal Sciences*, 24(2): 246-249. DOI: <https://www.doi.org/10.5713/ajas.2011.10060>
- Calik A, Emami NK, Schyns G, White MB, Walsh MC, Romero LF, and Dalloul RA (2022a). Influence of dietary vitamin E and selenium supplementation on broilers subjected to heat stress, Part II: oxidative stress, immune response, gut integrity, and intestinal microbiota. *Poultry Science*, 101(6): 101858. DOI: <https://www.doi.org/10.1016/j.psj.2022.101858>
- Calik A, Emami K, White B, Walsh C, Romero F, and Dalloul A (2022b). Influence of dietary vitamin E and selenium supplementation on broilers subjected to heat stress, Part I: Growth performance, body composition and intestinal nutrient transporters. *Poultry Science*, 101(6): 101857. DOI: <https://www.doi.org/10.1016/j.psj.2022.101857>
- Çavuşoğlu E, Petek M, Abdourhamane IM, Akkoc A, and Topal E (2018). Effects of different floor housing systems on the welfare of fast-growing broilers with an extended fattening period. *Archives Animal Breeding*, 61(1): 9-16. DOI: <https://www.doi.org/10.5194/aab-61-9-2018>
- Costa H, Vaz R, Silva M, Rodrigues K, Sousa L, Bezerra L, Ribeiro M, Barbosa A, Almeida J, and Oliveira M (2021). Performance and meat quality of broiler chickens reared on two different litter materials and at two stocking densities. *British Poultry Science*, 62(3): 396-403. DOI: <https://www.doi.org/10.1080/00071668.2020.1864810>
- Dalia A, Loh T, Sazili A, Jahromi M, and Samsudin A (2017). The effect of dietary bacterial organic selenium on growth performance, antioxidant capacity, and Selenoproteins gene expression in broiler chickens. *BMC Veterinary Research*, 13: 254. DOI: <https://www.doi.org/10.1186/s12917-017-1159-4>
- Dalia A, Loh T, Sazili A, Jahromi M, and Samsudin A (2018). Effects of vitamin E, inorganic selenium, bacterial organic selenium, and their combinations on immunity response in broiler chickens. *BMC Veterinary Research*, 14(1): 249. DOI: <https://www.doi.org/10.1186/s12917-018-1578-x>
- De Jong IC, Gunnink H, and Van Harn J (2014). Wet litter not only induces footpad dermatitis but also reduces overall welfare, technical performance, and carcass yield in broiler chickens. *Journal of Applied Poultry Research*, 23(1): 51-58. DOI: <https://www.doi.org/10.3382/japr.2013-00803>
- De Toledo TDS, Roll AAP, Rutz F, Dallmann HM, Dai Pra MA, Leite FPL, and Roll VFB (2020). An assessment of the impacts of litter treatments on the litter quality and broiler performance: A systematic review and meta-analysis. *Plos One*, 15(5): e0232853. DOI: <https://www.doi.org/10.1371/journal.pone.0232853>
- Deng S, Hu S, Xue J, Yang K, Zhuo R, Xiao Y, and Fang R (2022). Productive performance, serum antioxidant status, tissue selenium deposition, and gut health analysis of broiler chickens supplemented with selenium and probiotics—A pilot study. *Animals*, 12(9): 1086. DOI: <https://www.doi.org/10.3390/ani12091086>
- Duntas LH and Brenta G (2018). A renewed focus on the association between thyroid hormones and lipid metabolism. *Frontiers in Endocrinology*, 9: 511. DOI: <https://www.doi.org/10.3389/fendo.2018.00511>
- El-Maaty HA, Sherif S, Taha AE, Al-Otaibi AM, Othman SI, Allam A A, and Mahrose K (2023). Effects of housing systems and feed additive on growth, carcass traits, liver function, oxidative status, thyroid function, and immune parameters of broilers. *Poultry Science*, 102(12): 103121. DOI: <https://www.doi.org/10.1016/j.psj.2023.103121>
- Farghly MF, Mahrose KM, Cooper RG, Ullah Z, Rehman Z, and Ding C (2018). Sustainable floor type for managing turkey production in a hot climate. *Poultry Science*, 97(11): 3884-3890. DOI: <https://www.doi.org/10.3382/ps/pey280>
- Gangadoo S, Dinev I, Willson NL, Moore RJ, Chapman J, and Stanley D (2020). Nanoparticles of selenium as high bioavailable and non-toxic supplement alternatives for broiler chickens. *Environmental Science and Pollution Research*, 27: 16159-16166. DOI: <https://www.doi.org/10.1007/s11356-020-07962-7>

- Gao J, Lin H, Wang XJ, Song ZG, and Jiao HC (2010). Vitamin E supplementation alleviates the oxidative stress induced by dexamethasone treatment and improves meat quality in broiler chickens. *Poultry Science*, 89(2): 318-327. DOI: <https://www.doi.org/10.3382/ps.2009-00216>
- Ghanima MMA, Abd El-Hack ME, Othman SI, Taha AE, Allam AA, and Abdel-Moneim AME (2020). Impact of different rearing systems on growth, carcass traits, oxidative stress biomarkers, and humoral immunity of broilers exposed to heat stress. *Poultry Science*, 99(6): 3070-3078. DOI: <https://www.doi.org/10.1016/j.psj.2020.03.011>
- Ghasemi HA, Hajkhodadadi I, Hafizi M, Taherpour K, and Nazaran MH (2020). Effect of advanced chelate technology based trace minerals on growth performance, mineral digestibility, tibia characteristics, and antioxidant status in broiler chickens. *Nutrition & Metabolism*, 17: 94. DOI: <https://www.doi.org/10.1186/s12986-020-00520-5>
- Ghasemi HA, Kazemi-Bonchenari M, Khabtabadi-Farahani AH, and Khodaei Motlagh M (2013). The effect of feeding rations with different ratios of concentrate to alfalfa hay on blood hematological and biochemical parameters of farmed ostriches (*Struthio camelus*). *Tropical Animal Health and Production*, 45: 1635-1640. DOI: <https://www.doi.org/10.1007/s11250-013-0409-0>
- Gornall AG, Bardawill CJ, and David MM (1949). Determination of serum proteins by means of the biuret reaction. *Journal of Biological Chemistry*, 177(2): 751-766. Available at: <https://cir.nii.ac.jp/crid/1573105975550272896>
- Habibian M, Ghazi S, and Moeini MM (2016). Effects of dietary selenium and vitamin E on growth performance, meat yield, and selenium content and lipid oxidation of breast meat of broilers reared under heat stress. *Biological Trace Element Research*, 169: 142-152. DOI: <http://www.doi.org/10.1007/s12011-015-0404-6>
- Huang Y, Ren J, and Qu X (2019). Nanozymes: Classification, catalytic mechanisms, activity regulation, and applications. *Chemical Reviews*, 119(6): 4357-4412. DOI: <https://www.doi.org/10.1021/acs.chemrev.8b00672>
- Jo C and Ahn D (1998). Fluorometric analysis of 2-thiobarbituric acid reactive substances in turkey. *Poultry Science*, 77(3): 475-480. DOI: <https://www.doi.org/10.1093/ps/77.3.4>
- Kang B, Bansal M, and Mehta U (2000). Department of biophysics. Panjab University, Chandigarh-160014, India. *Biological Trace Element Research*, 77: 231-240. DOI: <https://www.doi.org/10.1385/BTER:77.3:231>
- Kazem SMH and Al-Khafaji FR (2023). Effect of addition different levels of organic, inorganic and nano-Se to the diet of broiler chickens (Ross 308) exposed to heat stress on characteristics blood cellular and some indicators of oxidative stress. *IOP Conference Series: Earth and Environmental Science*, 1262: 072001 Available at: <https://iopscience.iop.org/article/10.1088/1755-1315/1262/7/072001/meta>
- Lasheen HM, El-Gendi G, Iraqi MM, and El-Attrouny MM (2023). Effects of lighting colors and litter type on growth performance and physiological performance status of broilers. *Egyptian Poultry Science Journal*, 43(3): 489-504. DOI: <https://www.doi.org/10.21608/epsj.2023.319475>
- Li J, Miao Z, Tian W, Yang Y, Wang J, and Yang Y (2017). Effects of different rearing systems on growth, small intestinal morphology and selected indices of fermentation status in broilers. *Animal Science Journal*, 88(6): 900-908. DOI: <https://www.doi.org/10.1111/asj.12697>
- Lin SI, Wang CW, Tan Sr, Liang Y, Yao Hd, Zhang ZW, and Xu SW (2014). Selenium deficiency inhibits the conversion of thyroidal thyroxine (T₄) to triiodothyronine (T₃) in chicken thyroids. *Biological Trace Element Research*, 161: 263-271. DOI: <http://www.doi.org/10.1007/s12011-014-0083-8>
- Liu B, Xiong YL, Jiang J, Yu D, and Lin G (2021). Cellular antioxidant mechanism of selenium-enriched yeast diets in the protection of meat quality of heat-stressed hens. *Food Bioscience*, 39: 100798. DOI: <https://www.doi.org/10.1016/j.fbio.2020.100798>
- Mazur-Kuśnerek M, Antoszkiewicz Z, Lipiński K, Kaliniewicz J, and Kotlarczyk S (2019). The effect of polyphenols and vitamin E on the antioxidant status and meat quality of broiler chickens fed low-quality oil. *Archives Animal Breeding*, 62(1): 287-296. DOI: <https://www.doi.org/10.5194/aab-62-287-2019>
- Miller NJ, Rice-Evans C, Davies MJ, Gopinathan V, and Milner A (1993). A novel method for measuring antioxidant capacity and its application to monitoring the antioxidant status in premature neonates. *Clinical Science*, 84(4): 407-412. DOI: <https://www.doi.org/10.1042/cs0840407>
- National research council (NRC) (1994). Nutrient requirements of poultry. *Journal of Applied Poultry Research*, 3(1): 101. DOI: <https://www.doi.org/10.1093/japr/3.1.101>
- Negis Y, Aytan N, Özer N, Ogru E, Libinaki R, Gianello R, Azzi A, and Zingg JM (2006). The effect of tocopheryl phosphates on atherosclerosis progression in rabbits fed with a high cholesterol diet. *Archives of Biochemistry and Biophysics*, 450(1): 63-66. DOI: <https://www.doi.org/10.1016/j.abb.2006.02.027>
- Ognik K and Krauze M (2016). The potential for using enzymatic assays to assess the health of turkeys. *World's Poultry Science Journal*, 72(3): 535-550. DOI: <https://www.doi.org/10.1017/S0043933916000246>
- Okasha HM, El-Gendi GM, and El Garhy O (2021a). Effect of flock age, air pressure and litter types on carcass traits and meat quality of broiler chicks. *Annals of Agricultural Science, Moshtohor*, 59(2): 435-444. DOI: <http://www.doi.org/10.21608/ASSJM.2021.186382>
- Okasha HM, El-Gendi GM, and El Garhy O (2021b). Some factors affecting blood parameters of broiler chickens. *Annals of Agricultural Science, Moshtohor*, 59(2): 425-434. DOI: <http://www.doi.org/10.21608/ASSJM.2021.186382>
- Packer L, Weber SU, and Rimbach G (2001). Molecular Aspects of α -Tocotrienol Antioxidant Action and Cell Signalling. *The Journal of Nutrition*, 131(2): 369S-373S. DOI: <https://www.doi.org/10.1093/jn/131.2.369S>
- Paglia DE and Valentine WN (1967). Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *The Journal of Laboratory and Clinical Medicine*, 70(1): 158-169. DOI: <https://www.doi.org/10.5555/uri:pii:0022214367900765>
- Pepper CM and Dunlop MW (2021). Review of litter turning during a grow-out as a litter management practice to achieve dry and friable litter in poultry production. *Poultry Science*, 100(6): 101071. DOI: <https://www.doi.org/10.1016/j.psj.2021.101071>
- Perez-Carbajal C, Caldwell D, Farnell M, Stringfellow K, Pohl S, Casco G, Pro-Martinez A, and Ruiz-Feria CA (2010). Immune response of broiler chickens fed different levels of arginine and vitamin E to a coccidiosis vaccine and Eimeria challenge. *Poultry Science*, 89(9): 1870-1877. DOI: <https://www.doi.org/10.3382/ps.2010-00753>
- Rayman MP and Stranges S (2013). Epidemiology of selenium and type 2 diabetes: Can we make sense of it?. *Free Radical Biology and Medicine*, 65: 1557-1564. DOI: <https://www.doi.org/10.1016/j.freeradbiomed.2013.04.003>
- Renden J, Lien R, Oates S, and Bilgili S (1994). Plasma concentrations of corticosterone and thyroid hormones in broilers provided various lighting schedules. *Poultry Science*, 73(1): 186-193. DOI: <https://www.doi.org/10.3382/ps.0730186>
- Schomburg L (2012). Selenium, selenoproteins and the thyroid gland: Interactions in health and disease. *Nature Reviews Endocrinology*, 8(3): 160-171. DOI: <https://www.doi.org/10.1038/nrendo.2011.174>
- Selim N, Radwan N, Youssef S, Eldin TS, and Elwafa SA (2015). Effect of inclusion inorganic, organic or nano selenium forms in broiler diets on: 1-growth performance, carcass and meat characteristics. *International Journal of Poultry Science*, 14(3): 135-143. Available at: <https://www.doi.org/10.3923/ijps.2015.135.143>

- Şen G, Oktay MN, Evci Ş, Gökpinar S, and Şenol A (2023). The effect of using different litter materials in broiler rearing on performance, carcass yield, antioxidant status, some litter parameters, and coccidiosis oocysts. *Turkish Journal of Veterinary & Animal Sciences*, 47(5): 487-495. DOI: <https://www.doi.org/10.55730/1300-0128.4317>
- Shepherd E and Fairchild B (2010). Footpad dermatitis in poultry. *Poultry Science*, 89(10): 2043-2051. DOI: <https://www.doi.org/10.3382/ps.2010-00770>
- Sirois M (2019). Laboratory manual for laboratory procedures for veterinary technicians, 7th Edition. Elsevier Health Sciences, Available at: <https://www.uk.elsevierhealth.com/laboratory-manual-for-laboratory-procedures-for-veterinary-technicians-97803233595407.html>
- Surai PF, Kochish II, Fisinin VI, and Kidd MT (2019). Antioxidant defence systems and oxidative stress in poultry biology: An update. *Antioxidants*, 8(7): 235. Available at: <https://www.mdpi.com/2076-3921/8/7/235>
- Surai PF and Fisinin VI (2014). Selenium in poultry breeder nutrition: An update. *Animal Feed Science and Technology*, 191: 1-15. DOI: <https://www.doi.org/10.1016/j.anifeedsci.2014.02.005>
- Surai PF, Kochish II, Fisinin VI, and Velichko OA (2018). Selenium in poultry nutrition: From sodium selenite to organic selenium sources. *The Journal of Poultry Science*, 55(2): 79-93. DOI: <http://www.doi.org/10.2141/jpsa.0170132>
- Toth PP (2005). The good cholesterol high-density lipoprotein. *Circulation*, 111(5): e89-e91. DOI: <https://www.doi.org/10.1161/01.CIR.0000154555.07002>
- Traber MG and Stevens JF (2011). Vitamins C and E: Beneficial effects from a mechanistic perspective. *Free Radical Biology and Medicine*, 51(5): 1000-1013. DOI: <https://www.doi.org/10.1016/j.freeradbiomed.2011.05.017>
- Wang Q, Huang J, Zhang H, Lei X, Du Z, Xiao C, Chen S, and Ren F (2017). Selenium deficiency-induced apoptosis of chick embryonic vascular smooth muscle cells and correlations with 25 selenoproteins. *Biological Trace Element Research*, 176(2): 407-415. DOI: <https://www.doi.org/10.1007/s12011-016-0823-z>
- Wang Z, Kong L, Zhu L, Hu X, Su P, and Song Z (2021a). The mixed application of organic and inorganic selenium shows better effects on incubation and progeny parameters. *Poultry Science*, 100(2): 1132-1141. DOI: <https://www.doi.org/10.1016/j.psj.2020.10.037>
- Wang Y, Heng C, Zhou X, Cao G, Jiang L, Wang J, Li K, Wang D, and Zhan X (2021b). Supplemental *Bacillus subtilis* DSM 29784 and enzymes, alone or in combination, as alternatives for antibiotics to improve growth performance, digestive enzyme activity, anti-oxidative status, immune response and the intestinal barrier of broiler chickens. *British Journal of Nutrition*, 125(5): 494-507. DOI: <https://www.doi.org/10.1017/S0007114520002755>
- Wang Y, Ru YJ, Liu GH, Chang WH, Zhang S, Yan HJ, Zheng AJ, Lou RY, Liu ZY, and Cai HY (2015). Effects of different rearing systems on growth performance, nutrients digestibility, digestive organ weight, carcass traits, and energy utilization in male broiler chickens. *Livestock Science*, 176: 135-140. DOI: <https://www.doi.org/10.1016/j.livsci.2015.03.010>
- Wei F, Hu X, Xu B, Zhang M, Li S, Sun Q, and Lin P (2015). Ammonia concentration and relative humidity in poultry houses affect the immune response of broilers. *Genetics and Molecular Research*, 14(2): 3160-3169. DOI: <http://www.doi.org/10.4238/2015.April.10.27>
- Wickramasuriya SS, Park I, Lee Y, and Lillehoj HS (2023). Effect of dietary organic selenium on growth performance, gut health, and coccidiosis response in broiler chickens. *Animals*, 13(9): 1560. DOI: <https://www.doi.org/10.3390/ani13091560>
- Woolliams J, Woolliams G, Anderson P, and McMurray C (1983). Variation in the activities of glutathione peroxidase and superoxide dismutase and in the concentration of copper in the blood in various breed crosses of sheep. *Research in Veterinary Science*, 34(3): 253-256. DOI: [https://www.doi.org/10.1016/S0034-5288\(18\)32219-7](https://www.doi.org/10.1016/S0034-5288(18)32219-7)
- Yoon I, Werner TM, and Butler JM (2007). Effect of source and concentration of selenium on growth performance and selenium retention in broiler chickens. *Poultry Science*, 86(4): 727-730. DOI: <https://www.doi.org/10.1093/ps/86.4.727>
- Young J, Stagsted J, Jensen S, Karlsson A, and Henckel P (2003). Ascorbic acid, alpha-tocopherol, and oregano supplements reduce stress-induced deterioration of chicken meat quality. *Poultry Science*, 82(8): 1343-1351. DOI: <http://www.doi.org/10.1093/ps/82.8.1343>
- Zhang L, Wang Y, Zhou Y, Zheng L, Zhan X, and Pu Q (2014). Different sources of maternal selenium affect selenium retention, antioxidant status, and meat quality of 56-day-old offspring of broiler breeders. *Poultry Science*, 93(9): 2210-2219. DOI: <https://www.doi.org/10.3382/ps.2013-03605>
- Zhang Q, Qian ZY, Zhou PH, Zhou XL, Zhang DL, He N, Zhang J, Liu YH, and Gu Q (2018). Effects of oral selenium and magnesium co-supplementation on lipid metabolism, antioxidative status, histopathological lesions, and related gene expression in rats fed a high-fat diet. *Lipids in Health and Disease*, 17(1): 1-12. DOI: <https://www.doi.org/10.1186/s12944-018-0815-4>

Publisher's note: [Scenceline Publication](#) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025